



### **CHEMICAL KINETICS**

# **Rate of Chemical Reaction Fast Reactions** (Precipitation of silver chloride by mixing of aqueous solutions of silver nitrate and sodium chloride) Types of Reactions on the basis of speed Moderate Slow Reactions Reactions (Rusting of iron (Inversion of in the presence cane sugar and of air and hydrolysis of moisture) starch)

# Rate of Chemical Reaction

The **rate of reaction** is the change in concentration of a reactant or productin unit time.

• Alternatively, the rate of reaction can also be expressed as

The rate of decrease in concentration of any one of the reactants.

The rate of increase in concentration of any one of the products.

Consider a hypothetical reaction, assuming that the volume of the system remains constant.

$$R \rightarrow P$$

One mole of the reactant R produces one mole of the product P.

If [R]<sub>1</sub> and [P]<sub>1</sub> are the concentrations of R and P at time t<sub>1</sub> and [R]<sub>2</sub> and [P]<sub>2</sub> are their concentrations at time t2, then

$$\Delta \mathbf{t} = \mathbf{t}_2 - \mathbf{t}_1$$

$$\Delta \begin{bmatrix} \mathbf{R} \end{bmatrix} = \begin{bmatrix} \mathbf{R} \end{bmatrix}_2 - \begin{bmatrix} \mathbf{R} \end{bmatrix}_1$$

$$\Delta \begin{bmatrix} \mathbf{P} \end{bmatrix} = \begin{bmatrix} \mathbf{P} \end{bmatrix}_2 - \begin{bmatrix} \mathbf{P} \end{bmatrix}_1$$


The square brackets in the above expressions are used to express molar concentration.

Rate of disappearance of R = 
$$\frac{\text{Decrease in concentration of R}}{\text{Time taken}} = -\frac{\Delta[R]}{\Delta t}$$
 (1)

 $\Delta[R]$  is a negative quantity because the concentration of reactants is decreasing.

Rate of appearance of 
$$P = \frac{\text{Increase in concentration of P}}{\text{Time taken}} = + \frac{\Delta[P]}{\Delta t}$$
 (2)

Equations 1 and 2 represent the average rate of a reaction, rav. This average rate depends on the change in concentration of reactants or products and the timetaken for that change to occur.



### Units of Rate of a Reaction

- From Equations 1 and 2, it is clear that the units of rate are concentration time<sup>-1</sup>.
- For example, if concentration is in mol L<sup>-1</sup> and time is in seconds, then the units are mol L<sup>-1</sup>s<sup>-1</sup>.
- In gaseous reactions, the concentration of gases is expressed in terms of their partial pressures; hence, the units of the rate equation will be atm s<sup>-1</sup>.

### Instantaneous Rate of Reaction

• Consider the hydrolysis of butyl chloride (C<sub>4</sub>H<sub>9</sub>Cl).

$$C_4H_9CI+H_2O \rightarrow C_4H_9OH+HCI$$

• We have provided the concentrations over different intervals of time below.


Time (s <sup>-1</sup> )	0	50	100	150	200	300	400	700	800
Concentration (mol L <sup>-1</sup> )	0.100	0.0905	0.0820	0.0741	0.0671	0.0549	0.0439	0.0210	0.017

- We can determine the difference in concentration over different intervals of time, and thus, we determine the average rate by dividing  $\Delta[R]$  by  $\Delta t$ .
- It can be seen from experimental data that the average rate falls from  $1.90 \times 10^{-4}$  mol  $L^{-1}s^{-1}$  to  $0.4 \times 10^{-4}$  mol  $L^{-1}s^{-1}$ .
- However, the average rate cannot be used to predict the rate of reaction at a particular instant as itwould be constant for the time interval for which it is calculated.

- Hence, to express the rate at a particular moment of time, we determine the instantaneous rate.
- It is obtained when we consider the average rate at the smallest time interval, say dt, when  $\Delta t$  approaches zero.

Therefore, for an infinitesimally small dt, the instantaneous rate is given by

$$r_{inst} = -\frac{d[R]}{dt} = \frac{d[P]}{dt}$$



- By drawing the tangent at time ton either of the curves for the concentration of R versus time t or concentration of P versus time t and calculating the slope of the curve, we can determine the instantaneous rate of reaction.
- Hence, in this example, r<sub>inst</sub> at 600 s is calculated by plotting the graph of the concentration of butylchloride as against time t.
- A tangent is drawn on the curve at a point t = 600 s.

$$\therefore r_{\text{inst}} \text{ at } 600 \text{ s} = \left[ \frac{0.0165 - 0.037}{\left(800 - 400\right)} \right] \text{molL}^{-1} = 5.12 \times 10^{-5} \, \text{molL}^{-1} \text{s}^{-1}$$

$$\begin{array}{lll} \text{At } t = 250 \, s & r_{inst} = 1.22 \times 10^{-4} \, \text{mol} \, L^{-1} \, s^{-1} \\ t = 350 \, s & r_{inst} = 1.0 \times 10^{-4} \, \, \text{mol} \, L^{-1} \, s^{-1} \\ t = 450 \, s & r_{inst} = 6.4 \times 10^{-5} \, \, \text{mol} \, L^{-1} \, s^{-1} \end{array}$$

Now consider a reaction,

$$Hg(I) + CI_2(g) \rightarrow HgCI_2(s)$$

Here, the stoichiometric coefficients of the reactants and products are the same; hence, the rate of reaction is given as

Rate of reaction=
$$-\frac{\Delta[Hg]}{\Delta t} = -\frac{\Delta[Cl_2]}{\Delta t} = \frac{\Delta[HgCl_2]}{\Delta t}$$

Therefore, we can say that from the above equation that the rate of disappearance of any of the reactants is the same as the rate of appearance of the products.

Consider another reaction,

$$2HI(g) \rightarrow H_2(g) + I_2(g)$$

In this reaction, two moles of HI decompose to produce one mole each of H2 and I2, i.e. the stoichiometric coefficients of reactants or products are not equal to one; hence, we need to divide the rate of disappearance of any of the reactants or the rate of appearance of products by their respective stoichiometric coefficients.


Because the rate of consumption of HI is twice the rate of formation of H2 or I2, to make them equal, the term  $\Delta[HI]$  is divided by 2.

The rate of this reaction is given by


Rate of reaction = 
$$-\frac{1}{2} \frac{\Delta [HI]}{\Delta t} = \frac{\Delta [H_2]}{\Delta t} = \frac{\Delta [I_2]}{\Delta t}$$

For a gaseous reaction at constant temperature, concentration is directly proportional to the partial pressure of a species, and hence, the rate can be expressed as the rate of change in partial pressure of the reactant or the product.

# **Instantaneous Rate of Reaction**



# MIND MAP: LEARNING MADE SIMPLE CHAPTER - 4



# **Important Questions**

# **Multiple Choice questions-**

Question 1.A first order reaction has a half life period of 34.65 seconds. Its rate constant is

- (a)  $2 \times 10^{-2} \text{ s}^{-1}$
- (b)  $4 \times 10^{-4} \text{ s}^{-1}$
- (c) 20 s<sup>-1</sup>
- (d)  $2 \times 10^{-4} \text{ s}^{-1}$

Question 2.If a graph is plotted between In k and 1/T for the first order reaction, the slope of the straight line so obtained is given by

Question 3. The unit of rate constant for a zero order reaction is

- (a) mol L<sup>-1</sup>s<sup>-1</sup>
- (b) s<sup>-1</sup>
- (c) L mol<sup>-1</sup>s<sup>-1</sup>
- (d) L2mol<sup>-2</sup>s<sup>-1</sup>

Question 4.A catalyst increases the speed of a chemical reaction by

- (a) increasing activation energy
- (b) decreasing activation energy
- (c) increasing reactant energy
- (d) decreasing threshold energy

Question 5.The units of the rate constant for the second order reaction are:

- (a) mol<sup>-1</sup> litre s<sup>-1</sup>
- (b) mol litre<sup>-2</sup> s<sup>-1</sup>
- (c) s<sup>-1</sup>
- (d) mol litre<sup>-1</sup> s<sup>-1</sup>

Question 6. The value of k for a reaction is  $2.96 \times 10^{-30}$  s<sup>-1</sup>. What is the order of the reaction?

- (a) Zero
- (b) 3
- (c) 2
- (d) 1

Question 7.A reaction is found to be of second order with respect to concentration of carbon monoxide. If concentration of carbon monoxide is doubled, the rate of reaction will

- (a) triple
- (b) increase by a factor of 4
- (c) double
- (d) remain unchanged

Question 8. If the concentrations are expressed in mol litre-1 and time in s, then the units of rate constant for the first-order reactions are

- (a) mol litre<sup>-1</sup> s<sup>-1</sup>
- (b) mol<sup>-1</sup> litre s<sup>-1</sup>
- (c) s<sup>-1</sup>
- (d) mol<sup>2</sup> litre<sup>-2</sup> s<sup>-1</sup>

Question 9. The half life of a first order reaction having rate constant 200 s<sup>-1</sup> is

- (a)  $3.465 \times 10^{-2}$  s
- (b)  $3.465 \times 10^{-3}$  s
- (c)  $1.150 \times 10^{-2}$  S
- (d)  $1.150 \times 10^{-3}$  S

Question 10. The rate of a reaction is  $1.209 \times 10^{-4} L^2 \text{ mol}^{-2} \text{s}^4$ . The order of the reaction is:

- (a) zero
- (b) first
- (c) second
- (d) third

# **Very Short Question:**

Question 1. Is rate of reaction always constant?

Question 2. Can order of reaction be zero? Give example.

Question 3. What do you understand by rate law expression?

Question 4. Is it possible to determine or predict the rate law theoretically by merely looking at the equation?

Question 5. Define the term chemical kinetics?

Question 6. Define – Rate of reaction and the factors affecting the rate of reaction.

Question 7. What is average rate of a reaction? How is it determined?

Question 8. What are the units of rate of a reaction?

Question 9. Identify the reaction order for from each of the following rate constant –

- (a)  $k = 2.3 \times 10^{-5} L \ mol 1 \ s^{-1}$
- (b)  $k = 3.1 \times 10^{-4} s^{-1}$

Question 10. Consider the equation  $2NO(q)+2H_2(g)\rightarrow N_2(g)+2H_2O(g)$ 

The rate law for this equation is first order with respect to  $H_2$  and second order with respect to NO. write the rate law for this reaction.

# **Short Questions:**

Question 1 For the reaction A+B  $\rightarrow$  C+D, the rate of reaction doubles when the concentration of A doubles, provided the concentration of B is constant. To what order does A enter into the rate expression?

Question 2. . A chemical reaction 2A  $\stackrel{\Leftrightarrow}{\rightarrow}$  4B+C in gas phase occurs in a closed vessel. The concentration of B is found to be increased by  $5\times10^{-3}$  mole  $L^{-1}$  in 10 second. Calculate (i) the rate of appearance of B (ii) the rate of disappearance of A?

Question 3. For the following reactions, write the rate of reaction expression in terms of reactants and products?

$$4NH_3(g) + 5O_2(g)4 \rightarrow NO(g) + 6H_2O(g)$$
  
ii)  $2N_2O_5 2NO_2 + O_2$ 

Question 4. . The reaction  ${}^{2N_2O_5(g)}$  $\mathcal{O}_{2}(g)$  was studied and the following data were collected:

S.no (mol/L/min)	$[N_2O_0]$ maj $L^{-1}$	Rate of disappearance of $\left[N_2 O_5\right]$ (mol/L/min
1.	1.13>10-2	34×10 <sup>-5</sup>
2.	0.84 ×10 <sup>-2</sup>	25×10⁻⁵
3.	0.62×10 <sup>-2</sup>	18×10 <sup>-5</sup>

### **Determine**

- i) The order
- ii) The rate law.
- iii) Rate constant for the reaction.

Question 5. The following experimental data was collected for the reaction:

$$Cl_2(g) + 2NO(g) \rightarrow 2NOCl(g)$$

Trial	Intial conc. Of $Cl_2(mol/L)$	NO mol/L	Initial Rate,(mol/L/s)
1	0.10	0.010	1.2×10 <sup>-4</sup>

2	0.10	0.030	10.8×10 <sup>-4</sup>
3	0.20	0.030	21.6×10 <sup>-4</sup>

### Construct the rate equation for the reaction.

Question 6. Draw a graph for

- Concentration for order of reactant against time
- b) Log Ro/R against time for a first order reaction.

Question 7. In general it is observed that the rate of a chemical reaction doubles with every 10° rise in temperature. If this generalization holds for a reaction in the temperature 305K. range295K to what would be the **A** activation reaction?  $(R = 8.314 Jk^{-1} mol^{-1})$ 

Question 8. The rate constant for a reaction is  $5 \times 10^{9} \, s^{-1}$  at  $50^{9} \, C$  and  $4.5 \times 10^{7} \, s^{-1}$  at  $100^{9} \, C$ . Calculate the value of activation energy for the reaction  $R = 8.314 \text{ JK}^{-1} \text{mol}^{-1}$ ?

Question 9. Plot a graph showing variation of potential energy with reaction. coordinate?

Question 10. The conversion of molecules X to Y follows second order kinetics. If concentration of X is increased to three times how will it affect the rate of formation of Y?

# **Long Questions:**

Question 1. The rate of the chemical reaction doubles for an increase of 10 K in absolute temperature from 298 K. Calculate 12

Question 2. The activation energy for the reaction  $^{2HI_{(g)}} \rightarrow H_2 + I_{2(g)}$  is  $^{209.5\,\mathrm{k\,J\,mol^{-1}}}$  at 581K. Calculate the fraction of molecules of reactants having energy equal to or greater than activation energy?

Question 3. From the rate expression for the following reactions, determine their order of reaction and the dimensions of the rate constants.

) 
$${}^{3}NO(g) \rightarrow N_{2}O(g) Rate = k[NO]^{2}$$
  
(ii)  ${}^{4}N_{2}O_{2}(aq) + 3I^{-}(aq) + H^{+} \rightarrow 2H_{2}O(1) + I_{3}^{-} Rate = k[H_{2}O_{2}][I^{-}]$   
(iii)  ${}^{4}CHO(g) \rightarrow CH_{4}(g) + CO(g) Rate = k[CH_{3}CHO]^{3/2}$   
(iv)  ${}^{4}CH_{5}CI(g) \rightarrow C_{2}H_{2}(g) + HCI(g) Rate = k[C_{2}H_{5}CI]$ 

**Question 4**. The decomposition of  $^{\mathrm{NH}_3}$  on platinum surface is zero order reaction. What are

### **CHEMISTRY**

# CHEMICAL KINETICS

the rates of production of  $^{N_2}$  and  $^{H_2}$  if  $^k$  = 2.5×10<sup>-4</sup> mol<sup>-1</sup>L s<sup>-1</sup>?

**Question 5**. The decomposition of dimethyl ether leads to the formation of  $^{CH_4}$ ,  $^{H_2}$  and CO and the reaction rate is given by Rate =  $^{k[CH_3OCH_3]^{3/2}}$ . The rate of reaction is followed by increase in pressure in a closed vessel, so the rate can also be expressed in terms of the partial pressure of dimethyl ether, i.e.,  $^{Rate=k(P_{CH_3OCH_3})^{3/2}}$  If the pressure is measured in bar and time in minutes, then what are the units of rate and rate constants?

# **Assertion and Reason Questions:**

- 1. In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.
  - a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
  - b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
  - c) Assertion is correct statement but reason is wrong statement.
  - d) Assertion is wrong statement but reason is correct statement.

Assertion: The rate of reaction is always negative.

**Reason:** Minus sign used in expressing the rate shows that concentration of product is decreasing.

- **2.** In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.
  - a) Assertion and reason both are correct statements and reason is correct explanation for
  - b) Assertion and reason both are correct statements but reason is not correct explanation for assertionss.
  - c) Assertion is correct statement but reason is wrong statement.
  - d) Assertion is wrong statement but reason is correct statement.

**Assertion:** Kinetics explains the reaction mechanism.

Reason: Kinetics explains the formation of products.


# **Case Study Questions:**

1. In a reaction, the rates of disappearance of different reactants or rates of formation of different products may not be equal but rate of reaction at any instant of time has the same value expressed in terms of any reactant or product. Further, the rate of reaction may not depend upon the stoichiometric coefficients of the balanced chemical equation. The exact powers of molar concentrations of reactants on which rate depends are found experimentally and expressed in terms of 'order of reaction'. Each reaction has a characteristic rate constant depends upon temperature. The units of the rate constant depend upon the order of reaction.

The following questions are multiple choice questions. Choose the most appropriate answer:

- (i) The rate constant of a reaction is found to be  $3 \times 10^{-3}$  mol<sup>-2</sup>  $1^2$  sec<sup>-1</sup>. The order of the reaction is:
  - a) 0.5
  - b) 2
  - c) 3
  - d) 1
- (ii) In the reaction, A + 3B  $\rightarrow$  2C, the rate of formation of C is:
  - a) The same as rate of consumption of A.
  - b) The same as the rate of consumption of B.
  - c) Twice the rate of consumption of A.
  - d) 3232 times the rate of consumption of B.
- (iii) Rate of a reaction can be expressed by following rate expression, Rate =  $k[A]^2$  [B], if concentration of A is increased by 3 times and concentration of B is increased by 2 times, how many times rate of reaction increases?
  - a) 9 times
  - b) 27 times
  - c) 18 times
  - d) 8 times
- (iv) The rate of a certain reaction is given by, rate =  $k[H^+]^n$ . The rate increases 100 times when the pH changes from 3 to 1. The order (n) of the reaction is:
  - a) 2
  - b) 0
  - c) 1
  - d) 1.5

- (v) In a chemical reaction A + 2B → products, when concentration of A is doubled, rate of the reaction increases 4 times and when concentration of B alone is doubled rate continues to be the same. The order of the reaction is:
  - a) 1
  - b) 2
  - c) 3
  - d) 4
- **2.** The progress of the reaction,  $A \rightleftharpoons nB$  with time is represented in the following figure:



The following questions are multiple choice questions. Choose the most appropriate answer:

- (i) What is the value of n?
  - a) 1
  - b) 2
  - c) 3
  - d) 4
- (ii) Find the value of the equilibrium constant.
  - a) 0.6M
  - b) 1.2M
  - c) 0.3M
  - d) 2.4M
- (iii) The initial rate of conversion of A will be:

### **CHEMISTRY**

# CHEMICAL KINETICS

- a) 0.1 mol L<sup>-1</sup>hr<sup>-1</sup>
- b) 0.2 mol L<sup>-1</sup>hr<sup>-1</sup>
- c) 0.4 mol L<sup>-1</sup>hr<sup>-1</sup>
- d) 0.8 mol L<sup>-1</sup>hr<sup>-1</sup>
- (iv) For the reaction, if  $\frac{d[B]}{dt}=2\times 10^{-4}$  , value of  $\frac{d[A]}{dt}$  will be:
  - a)  $2 \times 10^{-4}$
  - b) 10<sup>-4</sup>
  - c)  $4 \times 10^{-4}$
  - d)  $0.5 \times 10^{-4}$
- (v) Which factor has no effect on rate of reaction?
  - a) Temperature.
  - b) Nature of reactant.
  - c) Concentration of reactant.
  - d) Molecularity.

# Answers key

# MCQ Answer:

- 1. Answer: (a)  $2 \times 10^{-2}$  s<sup>-1</sup>
- 2. Answer: : (a)  $-\frac{E_a}{R}$
- 3. Answer: (a) mol  $L^{-1}s^{-1}$
- 4. Answer: (b) decreasing activation energy
- 5. Answer: (a) mol<sup>-1</sup> litre s<sup>-1</sup>
- 6. Answer: (d) 1
- 7. Answer: (b) increase by a factor of 4
- 8. Answer: (c) s<sup>-1</sup>
- 9. Answer: (b)  $3.465 \times 10^{-3}$  s
- 10. Answer: (d) third

# **Very Short Answers:**

- 1. No. rate of a reaction is not always constant. It depends on many factors such as concentration, temperature etc.
- 2. Yes, decomposition of ammonia on a hot platinum surface is a zero order of reaction at high pressure
- 3. Answer: The rate law is the expression in which rate is given in terms of molar concentration of reactants with each term raised to some power, which may or may not be same as the stoichiometric coefficient of the reacting species in a balanced chemical equation.
- 4. Answer: No, the rate law cannot be predicted by merely looking at the balanced chemical equation but must be determined experimentally.
- 5. Answer: The branch of chemistry that deals with the study of reaction rates and their mechanis is called chemical Kinetics.
- 6. Answer: Rate of reaction can be defined as the change in concentration of a reactant or produc per unit time. Factors affecting the rate of reaction are temperature, concentration of reactants and catalyst.
- 7. Answer: Average rate of a reaction is defined as the change in concentration of a reactant or a product per unit time. It can be determined by dividing the change in concentration of reactant or product by the time interval

For the reaction:  $A \rightarrow B R av$ 

$$= \frac{-\Delta[A]}{\Delta t} = \frac{\Delta[B]}{\Delta t}$$

- 8. The units of rate of a reaction are Mol  $L^{-1}S^{-1}$  In gaseous reaction the unit of rate of reaction is atom.
- 9. a) Since the units of rate constant are  $Lmol^{-1}s^{-1}$  The reactions is of second order.
  - b) Since the units of rate constant are  $s^{-1}$ , The reaction is of first order
- 10. The rate law will be  $R = K[NO]^2[H_2]$

# **Short Answers:**

1. Answer

$$A+B \rightarrow C+D$$
  
 $Rate[A]^x$ 

Rate = 1 when A = 1 - - - 1Rate = 2 when A = 2 - - 2

Dividing equation 2) by 1)

$$\frac{2}{1}\alpha \frac{\left(2\right)^{x}}{1^{x}}$$

$$2^{1}\alpha \frac{\left(2\right)^{x}}{2^{x}}$$

 $\therefore x = 1$ 

The reaction is first order reaction.

### 2. Answer:

$$2A \rightarrow 4B$$

$$-\frac{1}{2} \frac{d[A]}{dt} = \frac{1}{4} \frac{d[B]}{dt} = \frac{d[C]}{dt}$$

Rate of disappearance of B i)

$$= \frac{5 \times 10^{-3}}{10 5} \text{mol } / \text{L}^{-1} = 5 \times 10^{-4} \text{ mol } \text{L}^{-1} \text{ s}^{-1}$$

$$= \frac{-\text{d}[A]}{dt} = \frac{2}{4} \frac{d[B]}{dt} = \frac{1}{2} \frac{d[B]}{dt}$$

$$\text{ii)} = \frac{1}{2} \times 5 \times 10^{-4} \text{ mol } \text{L}^{-1} \text{ s}^{-1} = 2.5 \times 10^{4} \text{ mol } \text{L}^{-1}$$

### 3. Answer:

In terms of reactant In terms of products
$$\begin{array}{c|c}
R_1 & \frac{1}{4} \frac{\Delta[NH_3]}{\Delta t} & R_3 = \frac{1}{4} \frac{\Delta[NO]}{\Delta t} \\
R_2 & \frac{-1}{5} \frac{\Delta[O_2]}{\Delta t} & R_4 = \frac{1}{6} \frac{\Delta[H_2O]}{\Delta t}
\end{array}$$

$$\frac{1}{4}R_{1} = \frac{1}{5}R_{2} = \frac{1}{4}R_{3} = \frac{1}{6}R_{4}$$

$$-\frac{1}{4}\frac{\Delta[NH_{3}]}{\Delta t} = -\frac{1}{5}\frac{\Delta[O_{2}]}{\Delta t} = \frac{1}{4}\frac{\Delta[NO]}{\Delta t} = \frac{1}{6}\frac{\Delta[H_{2}O]}{\Delta t}$$

II ) In terms of reactant In terms of product 
$$R1 = -\frac{\Delta[N_2O_5]}{\Delta t} \qquad R_2 = \frac{\Delta[NO_2]}{\Delta t}$$

$$R_3 = \frac{\Delta [O_2]}{\Delta t}$$

$$\frac{1}{2}R_1 = \frac{1}{2}R_2 = R_3 - \frac{\Delta[N_2O_5]}{\Delta t} = \frac{1}{2} \frac{\Delta[NO_2]}{\Delta t} = \frac{\Delta[O_2]}{\Delta t}$$

### 4. Answer:

Let the order of reaction be x

$$Rate = K[N_2O_5]^x$$

i) From the data -

$$18 \times 10^{-5} = (0.62 \times 10^{-2})^x$$
 \_\_\_\_\_\_3)

Dividing 1) by 2)

$$\frac{34 \times 10^{-5}}{25 \times 10^{-5}} = (\frac{1.13 \times 10^{-2}}{0.84 \times 10^{-2}})^{x}$$

$$(1.36) = {(1.35)}^x$$

X=1

The order of reaction with respect with respect to  $N_2O_5$  is 1

ii) Rate law  $R = K [N_2O_5]$ 

T8×107 mol/L/min Rate


iii) Rate constant,  $K = \overline{[N_2O_5]}$ 

### 5. Answer:

Order of NO is 2

Rate law =  $K[Cl_2]$ 

### 6. Answer



### 7. Answer:

$$T_1 = 295K T_2 = 305K$$

$$Ea = 2.303 R \left[ \frac{T_2 T_1}{T_2 - T_1} \right] \left[ \log \frac{k_2}{k_1} \right]$$

$$K_2 = 2k_1$$

$$E_a = 2.303 \times 8.314 \times \left[ \frac{305 \times 295}{305 - 295} \right] \log \frac{2k_1}{k_1}$$

$$= 2.303 \times 8.314 \times 8997.5 K \log 2$$

$$= 51855.2 \text{ J/mol} \quad (\log 2 = 0.3010)$$

### 8. Answer:


$$\log \frac{4.5 \times 10^7}{1.5 \times 10^7} = \frac{Ea}{2.303 \times 3.314} \left( \frac{373 - 323}{373 \times 323} \right)$$

$$\log 1.5 = \frac{Ea}{2.303 \times 3.314} \left( \frac{50}{373 \times 323} \right)$$

$$Ea = \left( \frac{2.303 \times 3.314 \times 373 \times 323}{50} \right) \times \log 1.5$$

$$= 22 \ KJ / mol$$

### 9. Answer:



Reaction coordinate ---->

### 10.Answer:

The reaction  $X \to Y$  follows second order kinetics. Therefore, the rate equation for this reaction will be:

Rate = 
$$k[X]^2$$
 (1)

Let  $X = a^{mol^{-1}}$ , then equation (1) can be written as:

$$Rate_1 = k.(a)_2$$

$$= ka_2$$

If the concentration of X is increased to three times, then X = Now, the rate equation will be:

Rate = 
$$k(3a)_2$$

$$=9(ka^2)$$

Hence, the rate of formation will increase by 9 times.

# **Long Answers:**

### 1. Answer:

It is given that T1 = 298 K

Therefore, 
$$T_2 = (298+10)K$$

We also know that the rate of the reaction doubles when temperature is increased by <sup>10°</sup>.

k and that of  $k_2 = 2k$ Therefore, let us take the value of k

Also, 
$$R = 8.314 \,\mathrm{J}\,\mathrm{K}^{-1}\,\mathrm{mol}^{-1}$$

Now, substituting these values in the equation:

$$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[ \frac{T_2 - T_1}{T_1 T_1} \right]$$

We get:

$$\log \frac{2k}{k} = \frac{E_*}{2.303 \times 8.314} \left[ \frac{10}{298 \times 308} \right]$$

$$\log 2 = \frac{E_a}{2.303 \times 8.314} \left[ \frac{10}{298 \times 308} \right]$$

$$E_{s} = \frac{2.303 \times 8.314 \times 298 \times 308 \times \log 2}{1.314 \times 298 \times 308 \times \log 2}$$

$$= 52897.78 \,\mathrm{J}\,\mathrm{mol}^{-1}$$

$$= 52.9 \,\mathrm{k \, J \, mol^{-1}}$$

### 2. Answer:

In the given case:

$$E_a = 209.5 \text{ kJ mol}^{-1} = 209500 \text{ J mol}^{-1}$$

$$T = 581 \text{ K}$$

$$R = 8.314 J K^{-1} mol^{-1}$$

Now, the fraction of molecules of reactants having energy equal to or greater than

activation energy is given as:

$$x=e^{-Ea/RT}$$

$$In^{x = -E_a/RT}$$

$$\log x = -\frac{E_a}{2.303 \, RT}$$

$$\log x = \frac{209500 \,\text{J} \,\text{mol}^{-1}}{2.303 \times 8.314 \,\text{J} \,\text{K}^{-1} \!\text{mol}^{-1} \times 581} = 18.8323$$

Now, x = Anti log (18.8323)

= Anti 
$$\log \frac{19.1677}{}$$

### 3. Answer:

(i) Given rate = 
$$k[NO]^2$$

Therefore, order of the reaction = 2

$$k = \frac{Rate}{[NO]^2}$$

Dimension of

$$= \frac{\text{mol } L^{-1} s^{-1}}{(\text{mol } L^{-1})^2}$$

$$= \frac{\text{m ol } L^{-1} s^{-1}}{\text{m ol }^{2} L^{-2}}$$

$$= L \operatorname{mol}^{-1} s^{-1}$$

(ii) Given rate = 
$$k[H_2O_2][I^-]$$

Therefore, order of the reaction = 2

$$k = \frac{Rate}{Rate}$$

Dimension of

$$= \frac{\text{mol } L^{-1} s^{-1}}{(\text{mol } L^{-1})(\text{mol } L^{-1})}$$

$$= L \, \text{mol}^{-1} \text{s}^{-1}$$

(iii) Given rate = 
$${}^{k}[CH_{3}CHO]^{3/2}$$

Therefore, order of reaction = 
$$\frac{3}{2}$$

$$k = \frac{\text{Rate}}{[\text{CH}_3\text{CHO}]^{\frac{3}{2}}}$$

$$\begin{split} &= \frac{\text{mol } L^{-1} \, s^{-1}}{(\text{mol } L^{-1})^{\frac{3}{2}}} \\ &= \frac{\text{mol } L^{-1} s^{-1}}{\text{mol}^{\frac{3}{2}} L^{-\frac{3}{2}}} \\ &= L^{\frac{1}{2}} \, \text{mol}^{-\frac{1}{2}} s^{-1} \end{split}$$

(iv) Given rate = 
$$k[C_2H_5Cl]$$
  
Therefore, order of the reaction = 1

Dimension of 
$$k = \frac{Rate}{[C_2H_5C1]}$$
$$= \frac{mol L^{-1}s^{-1}}{mol L^{-1}}$$
$$= s^{-1}$$

### 4. Answer:

The decomposition of  $\,^{\mathrm{NH}_{3}}$  on platinum surface is represented by the following equation.  $2NH_{3(g)} \xrightarrow{p_t} N_{2(g)} + 3H_{2(g)}$ 

Rate = 
$$-\frac{1}{2}\frac{d[NH_3]}{dt} = \frac{d[N_2]}{dt} = \frac{1}{3}\frac{d[H_2]}{dt}$$

However, it is given that the reaction is of zero order.

$$-\frac{1}{2}\frac{d[NH_3]}{dt} = \frac{d[N_2]}{dt} = \frac{1}{3}\frac{d[H_2]}{dt} = k$$

Therefore, = 
$$2.5 \times 10^{-4}$$
 mol L<sup>-1</sup>s

Therefore, the rate of production of  $N_2$  is

$$\frac{d[N_2]}{dt} = 2.5 \times 10^{-4} \text{mol } 10^{-5} \text{s}^{-1}$$

And, the rate of production of  $H_2$  is

$$\frac{d[H_2]}{dt} = 3 \times 2.5 \times 10^{-4} \text{ thol } L^{-1} \text{s}^{-1}$$
$$= 7.5 \times 10^{-4} \text{ mol } L^{-1} \text{s}^{-1}$$

### 5. Answer:

If pressure is measured in bar and time in minutes, then

$$Rate = k \left(P_{\text{CH}_{\text{1}}\text{OCH}_{\text{1}}}\right)^{\frac{3}{2}}$$

### **CHEMISTRY**

### CHEMICAL KINETICS

$$k = \frac{Rate}{\left(P_{\text{CH}_{i}} \infty H_{i}\right)^{\frac{3}{2}}}$$

Therefore, unit of rate constants  $= bar^{-1/2} min^{-1}$ 

$$(k) = \frac{bar \min^{-1}}{bar^{\frac{3}{2}}}$$

# **Assertion and Reason Answers:**

1. (d) Assertion is wrong statement but reason is correct statement.

### **Explanation:**

The rate reaction is never negative. Minus sign used in expressing the rate only shows that the concentration of the reactant is decreasing.

**2.** (a) Assertion and reason both are correct statements and reason is correct explanation for assertion.

### **Explanation:**

Kinetics deals with the reaction mechanism i.e., how the atoms rearrange themselves in the reactar molecules in a single step or a number of steps, finally leading to the product molecules.

# **Case Study Answers:**

### 1. Answer:

i. (c) 3

### **Explanation:**

Unit of k for  $n^{th}$  order =  $(mot \ L^{-1})^{1-n} sec^{-1}$ 

Here, 
$$k = 3 \times 10^{-3} \text{ mol}^{-2} L^2 \text{ sec}^{-1}$$

Unit of k = 
$$\text{mol}^{-2} L^2 \sec^{-1} \Rightarrow (\text{mol } L^{-1})^{-2} \sec^{-1}$$

Comparing (i) and (ii) we get,  $1 - n = -2 \Rightarrow n = 3$ 

II. (c) twice the rate of consumption of A.

### **Explanation:**

Rate 
$$=rac{d[A]}{dt}=-rac{1}{3}rac{d[B]}{dt}=rac{1}{2}rac{d[C]}{dt}$$

### iii. (c) 18 times

# **Explanation:**

Given, 
$$R_1 = k[A]^2 [B]$$

According to question, 
$$R_2 = k[3A]^2$$
 [2B]

= 
$$k \times 9 [A]^2 \times 2[B] = 18 \times k[A]^2 [B] = 18 R_1$$

iv. (c) 1

# **Explanation:**

Rate (r) = 
$$k[H^+]^n$$

When pH = 3; 
$$[H^+] = 10^{-3}$$

and when pH = 1; 
$$[H^+] = 10^{-1}$$

$$\therefore rac{r_1}{r_2} = rac{k(10^{-3})^n}{k(10^{-1})^n} \Rightarrow rac{1}{100} = \left(rac{10^{-3}}{10^{-1}}
ight)^n (\therefore r_2 = 100r_1)$$

$$\Rightarrow (10^{-2})^1 = (10^{-2})^n \Rightarrow n = 1$$

### V. (b) 2

### **Explanation:**

Let the order of reaction w.r.t. A is x and w.r.t. B is y.

$$r_1 = k[A]^x[B]^y$$

$$\mathbf{r}_2 = \mathbf{k}[2\mathbf{A}]^{\mathbf{x}}[\mathbf{B}]^{\mathbf{y}}$$

$$r_3=k[A]^x[2B]^y\\$$

$$\frac{r_1}{r_2} = \frac{k[A]^x[B]^y}{k[2A]^x[B]^y}$$

$$\Rightarrow \frac{1}{4} = \left(\frac{1}{2}\right)^{x} \Rightarrow \left(\frac{1}{2}\right)^{2} = \left(\frac{1}{2}\right)^{x} \Rightarrow x = 2$$

Similarly, 
$$\frac{r_1}{r_3} = \frac{k[A]^x[B]^y}{k[A]^x[2B]^y}$$

$$\Rightarrow 1 = \left(\frac{1}{2}\right)^y \Rightarrow \left(\frac{1}{2}\right)^0 = \left(\frac{1}{2}\right)^y \Rightarrow y = 0$$

Hence the rate law equation is

Rate =  $k[A]^2 [B]^0 \Rightarrow Order of reaction = 2$ 

# 2. Answer:

### i. (b) 2

### **Explanation:**

According to the figure,

in the given time of 4 hours ( 1 to 5) concentration of A falls from 0.5 to 0.3M,

while in the same time concentration of B increases from 0.2 to 0.6 M.

Decrease in concentration of A in 4 hours

$$= 0.5 - 0.3 = 0.2M$$

Increase in concentration of B in 4 hours

$$= 0.6 - 0.2 = 0.4M$$

Thus, increase in concentration of B in a given time is

twice the decrease in concentration of A. Thus, n = 2

ii. (b) 1.2M

### **Explanation:**

$$K = \frac{[B]^2}{[A]} = \frac{(0.6)^2}{0.3} = 1.2M$$

iii. (a) 0.1 mol L-1hr-1

### **Explanation:**

From 
$$t = 0$$
 to  $t = 1hr$ ,

For A, 
$$dx = 0.6 - 0.5 = 0.1 \text{ mol } L^{-1}$$

$$\therefore$$
 Initial rate of conversion of  $A=rac{\mathrm{d}x}{\mathrm{d}t}$ 

$$= \frac{0.1 \text{ mol L}^{-1}}{1 \text{hr}} = 0.1 \text{ mol L}^{-1} \text{hr}^{-1}$$

### **Explanation:**

$$A \rightleftharpoons 2B$$

$$-\frac{\mathrm{d[A]}}{\mathrm{dt}} = +\frac{1}{2} \frac{\mathrm{d[B]}}{\mathrm{dt}}$$

$$=\frac{1}{2}\times2\times10^{-4}=10^{-4}$$

# v. (d) Molecularity.

# **Explanation:**

The number of reacting species (atoms, ions or molecules) taking part in an elementary reaction is called molecularity and it has no influence on the rate of reaction.