

AMINES

Introduction

Amines are the derivatives of ammonia prepared by the replacement of one, two or all the three hydrogenatoms by alkyl and/or aryl groups.


Examples:

(i) CH₃-NH₂


Structure of Amines

- In amines, the nitrogen atom is trivalent and has an unshared pair of electrons. Hence the nitrogenorbitals are sp³ hybridised with pyramidal geometry.
- The three sp³ hybrid orbitals of nitrogen overlap with orbitals of hydrogen or carbon depending on thenature of the amines.
- The fourth orbital of nitrogen in all amines contains an unshared pair of electrons. It is due to the presence of unshared pair of electrons, the angle C-N-E is less than 109.5°.

For example: The bond angle for trimethyl amine is 108°.



Classification of Amines



R, R' and R" can be alkyl or aryl

Preparation of Amines

Reduction of Nitro Compounds

Nitro compounds on reduction with hydrogen gas in the presence of finely divided nickel, palladium orplatinum and also on reduction with metals in acidic medium give amines.

Ammonolysis

Alkyl halides or benzyl halide on reaction with an ethanolic solution of ammonia undergoes nucleophilic substitution reaction in which halogen atom is replaced by an amino (-NH₂) group. The process of cleavage of the C-X bond by ammonia molecule is known as ammonolysis.

$$\overset{\bullet}{N}H_3 + R - X \longrightarrow R\overset{+}{N}H_3\overset{-}{X}$$

Nucleophile

Substituted ammonium salt

The primary amine prepared behaves as a nucleophile and reacts with further alkyl halide to formsecondary, tertiary amines, and finally quaternary ammonium salt.

$$R - NH_2 \xrightarrow{RX} R_2 NH \xrightarrow{RX} R_3 N \xrightarrow{RX} R_4 \overset{+}{N} \overset{-}{X}$$

(1°) (2°) (3°) Quaternary animonium salt

The free amine can be obtained from the ammonium salt by treatment with a strong base.

$$RNH_3 \bar{X} + NaOH \longrightarrow R - NH_2 + H_2O + NaX$$

In this method, a mixture of primary, secondary and tertiary and also a quaternary ammonium salt. However a primary amine is prepared by taking large excess of ammonia. The order of reactivity of halides with amines is RI > RBr > RCI

Reduction of Nitriles

Nitriles on reducing with LiAlH₄ or catalytic hydrogenation produce primary amines.

$$CH_3C \equiv N \xrightarrow[Na(Hg)]{H_2/Ni} CH_3 - CH_2 - NH_2$$

Reduction of Amides

Amides on reducing with LiAlH₄ yield amines.

CH₃-C-NH₂
$$(i) \text{ LiAlH}_4 \rightarrow \text{CH}_3\text{-CH}_2\text{-NH}_2$$

Gabriel pthalimide synthesis

Pthalimide on reacting with ethanolic solution of KOH forms potassium salt of pthalimide which onheating with alkyl halide followed by alkaline hydrolysis yields the corresponding primary amine.

Hoffmann bromamide degradation reaction

In this method, primary amines are prepared by treating an amide with bromine in an aqueous orethanolic solution of NaOH.

The amine formed has one carbon atom less than the starting amide.

$$H_{3}C$$
 — C — NH_{2} + Br_{2} + $4NaOH$ — CH_{3} - NH_{2} + $Na_{2}CO_{3}$ + $2NaBr$ + $2H_{2}O$

Physical Properties of Amines

Solubility

Lower aliphatic amines are soluble in water because they can form a hydrogen bond with water. Solubility decreases with increase in molar mass of amines due to an increase in the size of thehydrophobic group.

Boiling points

Among the isomeric amines, primary and secondary amines have a high boiling point because they can form hydrogen bonds.

Tertiary amines cannot form hydrogen bonds due to the absence of a hydrogen atom for hydrogenbond formation.

Hence, the order of boiling points of isomeric amines is Primary > Secondary > Tertiary

Chemical Properties of Amines

a) Basic character of amines

- ❖ Amines have an unshared pair of electrons on the nitrogen atom due to which they act as a Lewisbase.
- ❖ The basic character of amines can be better understood in terms of their K_b and pK_b values.

$$R^{-}NH_{2} + H_{2}O = R - NH_{3} + OH^{-}$$

$$K = \frac{[R - NH_{3}][OH^{-}]}{[R - NH_{2}]H_{2}O]}$$

$$K[H_{2}O] = \frac{[R - NH_{3}][OH^{-}]}{[R - NH_{2}]}$$


$$K_{b} = \frac{[R - NH_{3}][OH^{-}]}{[R - NH_{2}]}$$

$$pK_{b} = -\log K_{b}$$

❖ Greater K_b value or smaller pK_b indicates that a base is strong.

b) Comparison of basic strength of aliphatic amines and ammonia

Aliphatic amines are stronger bases than ammonia due to the +I effect of alkyl groups, leading to highelectron density on the nitrogen atom.



c) Comparison of basic strength of primary, secondary and tertiary amines

- ❖ The order of basicity of amines in the gaseous phase follows the expected order on the basis ofthe +I effect:
- ❖ In aqueous solution, tertiary amines are less basic than either primary or secondary amines. This can be explained on the basis of the following factors:

Tertiary amine > Secondary amine > Primary amine > NH₃

Solvation effect

- ✓ Greater the stability of the substituted ammonium cation formed, stronger is the correspondingamine as a base.
- ✓ The tertiary ammonium ion is less hydrated than the secondary ammonium ion, which is less hydrated than the primary amine. Thus, tertiary amines have less tendency to form ammoniumion and consequently are least basic.

$$\begin{array}{c} OH_2 \\ H \\ \hline \\ N \\ \hline \\ OH_2 \\ \hline \\ OH_2 \\ \hline \\ 10 \\ \hline \\ Maximum \ hydration \\ \end{array}$$


✓ On the basis of the solvation effect, the order of basicity of aliphatic amines should be:Primary amine > Secondary amine > Tertiary amine NH₃

Steric factor

- ✓ As the crowding of the alkyl group increases from primary to tertiary, amine hindrance to hydrogen bonding increases which eventually decreases the basic strength. Thus, there is a subtle interplay of the inductive effect, solvation effect and steric hindrance of the alkyl group which decides the basic strength of alkyl amines in the aqueous state.
- ✓ When the alkyl group is small like CH₃, there is no steric hindrance to hydrogen bonding. In this case, the order of basicity in aqueous medium is: $(CH_3)_2NH > CH_3NH_2 > (CH_3)_3N > NH_3$
- ✓ When the alkyl group is the ethyl group, the group order of basicity in the aqueous medium is $(C_2H_5)_2NH > (C_2H_5)_3N > C_2H_5NH_2 > NH_3$

> Comparison of basic strength of aryl amines and alkanamines

- ✓ Generally, aryl amines are considerably less basic than alkyl amines. Example: Ethyl amine is more basic than aniline.
- ✓ In aniline, the -NH₂ group is directly attached to the benzene ring. Hence, the unshared pair of electrons on nitrogen is less available for protonation because of resonance.



- ✓ In the above resonating structures, there is a positive charge on the nitrogen atom making the lone pair less available for protonation. Hence, aniline is less basic than ethyl amine which hasno resonating structures.
- ✓ Less basicity of aniline can also be explained by comparing the relative stability of aniline and anilinium ion obtained by accepting a proton.
- ✓ Greater the number of resonating structures, greater is the stability of that species.
- ✓ Aniline is a resonance hybrid of five resonating structures, whereas anilinium ion

has only two resonating structures.

✓ Thus, aniline has less tendency to accept a proton to form the anilinium ion.

Effect of substituent on the basic character of amines

- ✓ An electron-donating or electron-releasing group (EDG) increases basic strength.
- ✓ An electron-withdrawing group (EWG) decreases basic strength.

EDG: releases electrons, stabilizes the cation and increases basic strength

EWG: withdraws electrons, destabilizes the cation and decreases basic strength

$$EDG = -CH_3, -OCH_3, -NH_2$$

$$EWG = -NO_2$$
, $-CN$, $-X$ (Ha $logen$)

Reaction of Amines

Acylation reaction

- > Aliphatic and aromatic primary and secondary amines (which contain replaceable hydrogen atoms) react with acid chlorides, anhydrides and esters to form substituted amide.
- ➤ The process of introducing an acyl group (R–CO–) into the molecule is called acylation.
- > The reaction is carried out in the presence of a stronger base than the amine, such

as pyridine, which removes HCl formed and shifts the equilibrium to the product side.

R-NH₂ + RCOCI
$$\xrightarrow{\text{Base}}$$
 RNHCOR + HCI
Acid Substitutd
amide

R'-NH₂ + R-C-O-C-R $\xrightarrow{\text{Base}}$ R'NHCOR + RCOOH

Acid Substituted amide

R'-NH₂ + R-C-O-C-R $\xrightarrow{\text{Substituted}}$ Substituted amide

R₂NH + RCOCI $\xrightarrow{\text{Base}}$ R₂NCOR + HCI

Important Note: Because tertiary amines do not contain a replaceable hydrogen atom, they do notundergo acylation.

Carbylamine reaction

- On heating aliphatic and aromatic primary amines with chloroform and ethanolic KOH they formisocyanides or carbylamines which have foul odour.
- Secondary and tertiary amines do not show this reaction.
- This reaction is used as a test for primary amines.

$$CH_3 - NH_2 + CHCI_3 + 3KOH \xrightarrow{Heat} CH_3 - NC + 3KCI + 3H_2O$$

Reaction with Nitrous acid

Primary aliphatic amines react with nitrous acid to form aliphatic diazonium salts. Being unstable diazonium salts liberate nitrogen gas quantitatively which is used in the estimation of amino acids and proteins.

$$\mathsf{CH_3} - \mathsf{NH_2} + \mathsf{HNO_3} \xrightarrow{\mathsf{NaNO_2} + \mathsf{HCI}} \\ \mathsf{[CH_3} - \overset{\mathsf{h}}{\mathsf{N_2}} \mathsf{CI}^{\scriptscriptstyle{-}}] \xrightarrow{\mathsf{H_2O}} \\ \mathsf{CH_3OH} + \mathsf{N_2} + \mathsf{HCI}$$

Aromatic amines on treating with nitrous acid at low temperatures to form diazonium salts which are used in the synthesis of a variety of aromatic compounds.

$$C_6H_5 - NH_2 \xrightarrow{NaNO_2 + HCI} C_6H_5 - N_2 CI^- + NaCI + 2H_2O$$

Secondary and tertiary amines react with nitrous acid in a different manner.

Reaction with arylsulphonyl chloride

Hinsberg's reagent or benzenesulphonyl chloride (C₆H₅SO₂Cl) reacts with primary amines and secondary amines to form sulphonamides.

Primary amine reacts with benzenesulphonyl chloride to form N-ethylbenzenesulphonyl amide.

N-ethylbenzenesulphonamide

The hydrogen bonded to nitrogen is strongly acidic due to the presence of strong electron withdrawing sulphonyl group and is hence soluble in alkali.

With secondary amine, N, N-diethyl-benzenesulphonamide is formed.

N,N-diethylbenzenesulphonamide

N, N-diethylbenzene sulphonamide does not contain any H atom attached to nitrogen atom so it is not acidic and is therefore insoluble in alkali.

> Tertiary amines do not react with benzenesulphonyl chloride.

Electrophilic substitution

Ortho- and para-positions to the -NH₂ group become centres of high electron density. So -NH₂ group is ortho and para directing and a powerful activating group.

(a)Bromination

Aniline reacts with bromine water at room temperature to give a white precipitate of 2, 4,6-tribromoaniline.

$$H_2$$
 H_2O H

Due to the high reactivity of aromatic amines, electrophilic substitution takes place at ortho- and para-positions.

For preparing monosubstituted aniline derivative, the -NH2 group is protected by acetylation with aceticanhydride then carrying out the desired substitution followed by the hydrolysis of the substituted amide to the substituted amine.

$$\frac{\text{OH-or H}^{+}}{\text{Pyridine}}$$

$$\frac{\text{OH-or H}^{+}}{\text{Br}_{2}}$$

$$\frac{\text{OH-or H}^{+}}{\text{Br}}$$

4-Bromoaniline

(b) **Nitration**

- Nitric acid is a nitrating agent plus a good oxidising agent. So direct oxidation of aromatic amines is not useful since it gives tarry oxidation products along with some nitro derivatives.
- In strong acidic medium, aniline is protonated to form the anilinium ion which is metal

directing. Hence besides the ortho and para derivatives, significant amount of meta derivative is also formed.

➤ However if we protect the −NH₂ group by acetylation reaction with acetic anhydride, the nitration reaction can be controlled and the p-nitro derivative derivative can be prepared as the major product.

(c) Sulphonation

Aniline on reacting with sulphuric acid forms anilinium hydrogen sulphate which on heating with sulphuric acid at 453-473K gives p-aminobenzene sulphonic acid as the major product.

Aniline does not undergo Friedel-Crafts reaction due to salt formation with Lewis acid aluminium chloride which is used as a catalyst. As a result, nitrogen of aniline acquires positive charge and hence acts as a strong deactivating group for further reaction.

Diazonium Salts

- Diazonium salts have the general formula

 RN2
 - ✓ Where R = Aryl group
 - \checkmark X⁻ion = Cl⁻ Br⁻, HSO₄, BF₄ etc.
- A suffix diazonium is added to the parent hydrocarbon from which they are formed followed by thename of the anion.
 - ✓ Anion = chloride, hydrogensulphate, etc.
 - ✓ Diazonium group = N^{+}_{2}
- Primary aliphatic amines form highly unstable alkyldiazonium salts whereas primary aromatic aminesform arenediazonium salts which are stable for a short time in a solution at low temperatures.
- The stability of arenediazonium ion is explained on the basis of resonance.

Preparation of Diazonium Salts

- Benzenediazonium chloride is prepared by the action of aniline with nitrous acid at 273-
- The conversion of primary aromatic amines into diazonium salts is known as diazotisation.

$$C_6H_5NH_2 + NaNO_2 + 2HCI \xrightarrow{273-278K} C_6H_5 \xrightarrow{1}_{0} CI^- + NaCI + 2H_2O$$

Physical Properties


- Benzenediazonium chloride is a colourless crystalline solid,
- It is readily soluble in water and is stable in cold but reacts with warm water.
- It decomposes easily in the dry state.
- Benzenediazonium fluoborate is water insoluble and stable at room temperature.

Chemical Reactions


Reactions involving displacement of Nitrogen

Replacement by halide or cyanide ion:

This reaction is called Sandmeyer reaction in which nucleophiles like Cl⁻,Br⁻ and CN⁻ can be easilyintroduced in the benzene ring in the presence of Cu(I) ion.



Alternatively, chlorine or bromine can also be introduced in the benzene ring by treating the diazonium salt solution with corresponding halogen acid in the presence of Cu powder. This isreferred as Gatterman reaction.



> Replacement by iodide ion:

Iodobenzene is formed on treating diazonium salt solution with potassium iodide.

$$ArN_2X$$
 + KI \longrightarrow ArI + KCI + N_2

> Replacement by fluoride ion:

Arenediazonium chloride on treating with fluoboric acid gives a precipitate of arene diazoniumfluoroborate which on heating decomposes to give aryl fluoride.

$$ArN_2CI + HBF_4 \longrightarrow Ar-N_2BF_4 \longrightarrow Ar-F + BF_3 + N_2$$

Replacement by H:

Hypophosphorus acid or ethanol are mild reducing agents and reduce diazonium salts to arenesand themselves get oxidised to phosphorus acid and ethanal respectively.

$$ArN_2CI + H_3PO_3 + H_2O$$
 $ArH + N_2 + H_3PO_3 + HCI$
 $ArN_2CI + CH_3CH_2OH$ $ArH + N_2 + CH_3CHO + HCI$

> Replacement by hydroxyl group:

Diazonium salt solution gets hydrolysed to phenol when the temperature is allowed to rise up to 283K.

$$ArN_2CI + H_2O$$
 \longrightarrow ArOH + N_2 + HCI

➤ Replacement by –NO₂ group:

On heating diazonium fluoroborate with aqueous sodium nitrite solution in the presence of copper, the diazonium group is replaced by -NO₂ group.

$$+ HBF_4 \longrightarrow NaNO_2 + N_2 + NaBF_4$$

Reactions involving retention of diazo group coupling reactions

- ✓ Benzene diazonium chloride reacts with phenol in which the phenol at its para position is coupled with the diazonium salt to form orange colour dye called phydroxyazobenzene.
- ✓ The reaction of diazonium salt with aniline gives yellow dye p-aminoazobenzene.
- ✓ The reaction is known as coupling reaction and it is an example of electrophilic substitutionreaction.

$$N = NC\bar{l} + H$$
 $NH_2 = H^+$


p-aminoazobenzene (yellow dye)

p-hydroxyazobenzene (orange dye)

Importance of Diazonium Salts in Synthesis of Organic Compounds

- Diazonium salts are very good intermediates for introducing –F, –Cl, –Br, –l, –CN, –OH, NO₂ groupsinto the aromatic ring.
- Direct halogenation method cannot be used for preparing aryl fluorides and iodides.
- Cyanobenzene can be easily prepared from diazonium salt.
- Thus the replacement of diazo group by other groups is useful in preparing substituted aromatic compounds which cannot be prepared by direct substitution in benzene or substituted benzene.

MIND MAP: LEARNING MADE SIMPLE CHAPTER - 13



Important Questions

Multiple Choice questions-

Question 1. Which of the following does not react with Hinsberg reagent?

- (a) Ethylamine
- (b) (CH₃)₂NH
- (c) $(CH_3)_3N$
- (d) Propan-2-amine

$$C_2H_5NH_2 \xrightarrow{NaNO_2/HCI} X \xrightarrow{P/Br_2} Y \xrightarrow{NH_3} Z$$

Question 2.

in above sequence, Z is

- (a) Cyanoethane
- (b) Ethanamide
- (c) Methanamine
- (d) Ethanamine

Question 3. Oxidation of aniline with K₂Cr₂O₇/H₂SO₄ gives

- (a) phenylhydroxylamine
- (b) p-benzoquinone
- (c) nitrosobenzene
- (d) nitrobenzene

Question 4. Which of the following amines can exhibit enantiomerism?

- (a) Benzeamine
- (b) 2-Butanamine
- (c) 2-Propanamine
- (d) 2-Methyl-propanamine.

Question 5. Which of the following: when heated with a mixture of ethanmine and alcoholic potas gives ethyl isocyanide?

- (a) 2-chloropropane
- (b) 2,2-dichloropropane
- (c) trichloromethane
- (d) tetrachloromethane

Question 6. Which of the following pair of species will yield carbylamine?

- (a) CH₃CH₂Br and KCN
- (b) CH₃CH₂Br and NH₃ (excess)
- (c) CH₃CH₂Br and AgCN
- (d) CH₃CH₂NH₂ and HCHO

Question 7. Which one of the following methods is neither meant for the synthesis nor for separation of amines?

- (a) Hinsberg method
- (b) Hoffmann method
- (c) Wurtz reaction
- (d) Curticus reaction

Question 8. C₆H₅CONHCH₃ can be converted into C₆H₅CH₂NHCH₃ by

- (a) NaBH₄
- (b) H₂-Pd/C
- (c) LiAlH₄
- (d) Zn-Hg/HCl

Question 9. $C_6H_5N^+_2Cl^- + CuCN \rightarrow C_6H_5CN + N_2 + CuCl$. The above chemical reaction is associated w which of the following name:

- (a) Balz Schiemen
- (b) Gattermann
- (c) Shimonini
- (d) Sandmeyer.

Question 10. The reaction of aniline with benzoyl chloride gives

- (a) Benzoin
- (b) Benzanilide
- (c) Benzalaniline
- (d) Benzamide

Very Short Questions-

- 1. For an amine RNH₂, write an expression to indicate its basic strength.
- 2. Give one use of quaternary ammonium salts.
- 3. Give one example of Hoffmann Bromamide reaction
- 4. Distinguish between ethylamine and aniline.
- 5. The IUPAC name of secondary amine having lowest molecular mass is ______.
- 6. Give an example of diazotization
- 7. Write one use of diazonium salt
- 8. How can the reactivity of aromatic amines be controlled?
- 9. Give one use of tertiary amines.
- 10. Name a reagent which can distinguish between primary, secondary and tertiary amine

Short Questions--

- 1. It is difficult to prepare pure amines by ammonolysis of alkylhalides.
- 2. Amines have higher boiling points than hydrocarbons of similar molecular mass.
- 3. Aniline is weaker base than cyclohexylamine.
- 4. Methylamine is a stronger base than aniline.

- 5. Before nitration, aniline is converted to acetanilide.
- 6. It is easier to brominate aniline as compared to benzene.
- 7. Reduction of nitro compound to aniline using iron scrap and HCl is preferred.
- 8. Aromatic amines cannot be prepared by Gabriel Phthalimide synthesis.
- 9. During acylation of amines, pyridine is added.
- 10. Aniline does not undergo Friedel Craft's reaction.

Long Questions-

- 1. (i) Write structures of different isomeric amines corresponding to the molecular formula, $^{\rm C_4H_{11}N}$
- (ii) Write IUPAC names of all the isomers.
- (iii) What type of isomerism is exhibited by different pairs of amines?
- 2. Arrange the following in increasing order of their basic strength:
- (i) C₂H₅NH₂,C₆H₅NH₂,NH₃,C₆H₅CH₂NH₂ and (C₂H₅)₂ NH
- (ii) $C_2H_5NH_2$, $(C_2H_5)_2$ NH, $(C_2H_5)_3$ N, $C_6H_5NH_2$
- (iii) CH₃NH₂,(CH₃)₂NH,(CH₃)₃N,C₆H₅NH₂,C₆H₅CH₂NH₂
- 3. Write structures of different isomers corresponding to the molecular formula, ${}^{C_3H_{\mathfrak{g}}N}$. Write IUPAC names of the isomers which will liberate nitrogen gas on treatment with nitrous acid.
- 4. Write IUPAC names of the following compounds and classify them into primary, secondary and tertiary amines.
- (i) $(CH_3)_2 CHNH_2$ (ii) $CH_3 (CH_2)_2 NH_3$
- (iii) CH₃NHCH(CH₃)₂ (iv) (CH₃)₃ CNH₂
- (v) $C_6H_5NHCH_3$ (vi) $(CH_3CH_2)_2NCH_3$ (vii) $m-BrC_6H_4NH_2$
- 5. Give one chemical test to distinguish between the following pairs of compounds.
- (i) Methylamine and dimethylamine
- (ii) Secondary and tertiary amines
- (iii) Ethylamine and aniline
- (iv) Aniline and benzylamine
- (v) Aniline and N-methylaniline.
- 6. Account for the following:
- (i) pKb of aniline is more than that of methylamine.
- (ii) Ethylamine is soluble in water whereas aniline is not.
- (iii) Methylamine in water reacts with ferric chloride to precipitate hydrated ferric oxide.
- (iv) Although amino group is o, p- directing in aromatic electrophilic substitution reactions, aniline on nitration gives a substantial amount of m-nitroaniline.

- (v) Aniline does not undergo Friedel-Crafts reaction.
- (vi) Diazonium salts of aromatic amines are more stable than those of aliphatic amines.
- (vii) Gabriel phthalimide synthesis is preferred for synthesising primary amines.
- 7. Arrange the following:
- (i) In decreasing order of the pKbvalues:

$$C_2H_{\epsilon}NH_2$$
, $C_6H_{\epsilon}NHCH_3$, $(C_2H_{\epsilon})_2NH_{and}C_6H_{\epsilon}NH_2$

(ii) In increasing order of basic strength:

$$C_6H_5NH_2$$
, $C_6H_5N(CH_3)_2$, $(C_2H_5)_2NH_{and}CH_3NH_2$

- (iii) In increasing order of basic strength:
- (a) Aniline, p-nitroaniline and p-toluidine
- (b) C₆H_ENH₂, C₆H_ENHCH₃, C₆H_ECH₂NH₂.
- (iv) In decreasing order of basic strength in gas phase:

$$C_2H_{\epsilon}NH_2$$
 , (C2H5)2NH, $(C_2H_{\epsilon})_3N_2$ and NH_3

(v) In increasing order of boiling point:

$$C_2H_5OH_1$$
 $(CH_3)_2NH_1$ $C_2H_5NH_2$

(vi) In increasing order of solubility in water:

$$C_6H_5NH_2$$
 $(C_2H_5)_2NH$ $C_6H_5NH_2$

- 8. How will you convert:
- (i) Ethanoic acid into methanamine
- (ii) Hexanenitrile into 1-aminopentane
- (iii) Methanol to ethanoic acid
- (iv) Ethanamine into methanamine
- (v) Ethanoic acid into propanoic acid
- (vi) Methanamine into ethanamine
- (vii) Nitromethane into dimethylamine
- (viii) Propanoic acid into ethanoic acid

Assertion and Reason Questions-

- 1. In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.
 - a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
 - b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
 - c) Assertion is correct statement but reason is wrong statement.
 - d) Assertion is wrong statement but reason is correct statement.

Assertion: Ortho substituted anilines are usually weaker bases than anilines.

Reason: This is due to ortho effect.

2. In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.

- a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
- b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
- c) Assertion is correct statement but reason is wrong statement.
- d) Assertion is wrong statement but reason is correct statement.

Assertion: In strongly acidic solutions, aniline becomes more reactive towards electrophilic reagents.

Reason: The amino group being completely protonated in strongly acidic solution, the lone pair of electrons on the nitrogen is no longer available for resonance.

Case Study Questions-

1. Read the passage given below and answer the following questions:

Amines are alkyl or aryl derivatives of ammonia formed by replacement of one or more hydrogen atoms. Alkyl derivatives are called aliphatic amines and aryl derivatives are known as aromatic amines. The presence of aromatic amines can be identified by performing dye test. Aniline is the simplest example of aromatic amine. It undergoeselectrophilic substitution reactions in which $-NH_2$ group strongly activates the aromatic ring through delocalisation of lone pair of electrons of N-atom. Aniline undergoes electrophilic substitution reactions. Ortho and para positions to the $-NH_2$ group become centres of high electrons density. Thus, $-NH_2$ group is ortho and para-directing and powerful activating group.

The following questions are multiple choice questions. Choose the most appropriate answer:

- (i) Cyclohexylamine and aniline can be distinguished by:
 - a) Hinsberg test.
 - b) carbylamine test.
 - c) Lassaigne test.
 - d) azo dye test.
- (ii) Which of the following compounds gives dye test?

- a) Aniline.
- b) Methyl amine.
- c) Diphenyl amine.
- d) Ethyl amine.
- (iii) Aniline when acetylated, the major product on nitration followed by alkaline hydrolysis gives:
 - a) Acetanilide.
 - b) o-nitroacetanitide.
 - c) p-nitroaniline.
 - d) m-nitroanitine.
- (iv) Oxidation of aniline with manganese dioxide and sulphuric acid produces:
 - a) Phenylhydroxylamine.
 - b) Nitrobenzene.
 - c) p-benzoquinone.
 - d) Phenol.
- (v) Aniline when treated with cone. HNO₃ and H₂SO₄ gives:
 - a) p-phenylenediamine.
 - b) m-nitroaniline.
 - c) p-benzoquinone.
 - d) Nitrobenzene.
- 2. Read the passage given below and answer the following questions:

The amines are basic in nature due to the presence of a lone pair of electron on N-atom of the -NH₂ group, which it can donate to electron deficient compounds. Aliphatic amines are stronger bases than NH₃ because of the +I effect of the alkyl groups. Greater the number of alkyl groups attached to N-atom, higher is the electron density on it and more will be the basicity. Thus, the order of basic nature of amines is expected to be $3^{\circ} > 2^{\circ} > 1^{\circ}$, however the observed order is $2^{\circ} > 1^{\circ} > 3^{\circ}$. This is explained on the basis of crowding on N-atom of the amine by alkyl groups which hinders the approach and bonding by a proton, consequently, the electron pair which is present on N is unavailable for donation and hence 3° amines are the weakest bases.

Aromatic amines are weaker bases than ammonia and aliphatic amines. Electron-donating groups such as $-CH_3$, $-OCH_3$, etc. increase the basicity while electron-withdrawing substitutes such as $-NO_2$, -CN, halogens, etc. decrease the basicity of amines. The effect of these substituents is more at p than at m-positions.

The following questions are multiple choice questions. Choose the most appropriate answer:

- (i) Which one of the following is the strongest base in aqueous solution?
 - a) Methyl amine.
 - b) Tri methyl amine.
 - c) Aniline.
 - d) Dimethyl amine.
- (ii) Which order ofbasicity is correct?
 - a) Aniline > m-toluidine > o-toluidine
 - b) Aniline> o-toluidine > m-toluidine
 - c) o-toluidine > aniline > m-toluidine
 - d) o-toluidine < aniline < m-toluidine
- (iii) What is the decreasing order of basicity of primary, secondary and tertiary ethylamines and NH₃?
 - a) $NH_3 > C_2H_5NH_2 > (C_2H_5)_2NH > (C_2H_5)_3N$
 - b) $(C_2H_5)_3N > (C_2H_5)_2NH > C_2H_5NH_2 > NH_3$
 - c) $(C_2H_5)_2NH > C_2H_5NH_2 > (C_2H_5)_3N > NH_3$
 - d) $(C_2H_5)_2NH > (C_2H_5)_3N > C_2H_5NH_2 > NH_3$
- (iv) The order of basic strength among the following amines in benzene solution is:
 - a) $CH_3NH_2 > (CH_3)_3N > (CH_3)_2NH$
 - b) $(CH_3)_3N > (CH_3)_2NH > CH_3NH_2$
 - c) $CH_3NH_2 > (CH_3)_2NH > (CH_3)_3N$
 - d) $(CH_3)_3N > CH_3NH_2 > (CH_3)_2NH$
- (v) Choose the correct statement.
 - a) Methylamine is slightly acidic.
 - b) Methylamine is less basic than ammonia.
 - c) Methylamine is a stronger base than ammonia.
 - d) Methylamine forms salts with alkalies.

MCQ Answers-

- 1. Answer: (c) (CH₃)₃N
- 2. Answer: (d) Ethanamine

SCIENCE

AMINES


- 3. Answer: (b) p-benzoquinone
- 4. Answer: (b) 2-Butanamine
- 5. Answer: (c) trichloromethane
- 6. Answer: (c) CH₃CH₂Br and AgCN
- 7. Answer: (c) Wurtz reaction
- 8. Answer: (c) LiAlH₄
- 9. Answer: (d) Sandmeyer.
- 10. Answer: (b) Benzanilide

Very Short Answers-

Ans 1.

$$RNH_2 + H_2O \implies RNH_3^+ + OH^-$$


$$K_b = \frac{[RNH_3^+] [OH^-]}{[RNH_3]}$$



Ans 3. In Hoffmann – Bromamide reaction an acid amide is reacted with Bromine in presence of a base to give a primary amine having one carbon less than the starting amide.

$$RCONH_2 + Br_2 + 4 \text{ NaOH } \rightarrow RNH_2 + Na_3 CO_3 + 2 \text{ NaBr} + 2 H_2O$$

Ans 4. Ethylamine and aniline can be distinguished by azodye test. On treating aniline with benzen diazonium salt, orange or red coloured azodye is formed which is not formed with ethyl amine.



Ans 5. N- Methylmethanamine.

Ans 6. During diazotization benzene diazonium chloride is prepared by the reaction of aniline with nitrous acid at 273 – 278 K

- Ans 7. Diazonium salts are used in preparation of substituted aromatic compounds.
- Ans 8. The reactivity of aromatic amines can be controlled by acylation.
- **Ans 9.**Tertiary amines like trimethylamine are used as insect attractants.
- **Ans 10.** P- Toluenesulphonyl chloride Hinsberg reagent can be used as a distinguishing reagent for primary, secondary and tertiary amines.

Short Answers-

Ans 1. The process of ammonolysis yields a mixture of primary, secondary, tertiary amines and quaternary salts. The separation of this amines is a very complicated process and difficult. Therefore it is difficult to prepare pure amines by ammonolysis of alkyl halides.


$$Rx \xrightarrow{NH_3} RNH_2 \xrightarrow{Rx} R_2NH \xrightarrow{Rx} R_3N \xrightarrow{Rx} [R_3N]^+x^-$$

Ans 2. Amines have higher boiling points than hydrocarbons of comparable molecular mass due to the presence of intermolecular hydrogen bond in amines which is absent in hydrocarbons. Therefore, amines exist as associated molecules and have higher boiling points.


In alcohols and carboxylic acids, the electro- negativity of oxygen is more than nitrogen of amines. Therefore the hydrogen bonds of alcohols and acids are stronger than in amines and alcohols & carboxylic acids have higher boiling points.

Ans 3. As a result of resonance in aniline; the lone pair on nitrogen delocalized over the benzene ri and is less available for protonation than in cyclohexyl amine which does not undergo resonance. Resonating structure of aniline —

Ans 4. Due to electron donating nature of CH_3 , group, electron pair. Availability at N- atom in meth amine is much higher than that in aniline; in aniline the benzene ring decreases the electron densit at N- atom in aniline. Thus CH_3NH_2 is a stronger base than aniline.



Ans 5. Aniline is very much susceptible to nitration and nitric acid is a strong oxidizing agent. Therefore to avoid oxidation of aniline, ${}^{-NH_2}$ group is protected by its acetylation to acetanilide which undergo nitration smoothly without any oxidation.



Ans 6. In aniline, due to +R effect of $^{-NH_2}$ group the benzene ring gets activated to a large extent a it becomes easier to brominates aniline as compared to benzene.

Ans 7. For reduction of nitro compounds to aniline, iron scrap and HCl is preferred because FeCl_2 formed gets hydrolysed to release HCl during the reaction & therefore only a small amount of HCl required to initiate the reaction.

Ans 8. Aromatic amines cannot be prepared by Gabriel pythalimide synthesis as aryl halides do not undergo nucleophilic substitution with the anion formed by pythalimide.

Ans 9. Acylation of amines is carried out in presence of pyridine or another base stronger than amines as it removes HCl so formed and shifts the equilibrium in forward direction.

$$C_2H_5 NH_2 + CH_3 COCI$$
Base
 $C_2H_5 NHCO CH_3 + HCI$

Ans 10. During Fridel Craft's reaction, aniline forms salt with aluminum chloride, the catalyst of reaction due to which nitrogen acquires a positive charge and acts as a strong deactivating group further reaction.

Long Answers-

Ans 1. (i), (ii) The structures and their IUPAC names of different isomeric amines corresponding to the molecular formula, $^{C_4H_{11}N}$ are given below:

Butanamine (1°)

2-Methylpropan-2-amine (1°)

N-Methylpropanamine (2°)

N-Methylpropan-2-amine

(iii) The pairs (a) and (b) and (e) and (g) exhibit position isomerism.

The pairs (a) and (c); (a) and (d); (b) and (c); (b) and (d) exhibit chain isomerism.

The pairs (e) and (f) and (g) exhibit metamerism.

All primary amines exhibit functional isomerism with secondary and tertiary amines and vice-versa

Ans 2. (i) Considering the inductive effect of alkyl groups, NH_3,C_2H_5NH_2 and $^{(C_2H_5)_2NH}$ can be arranged in the increasing order of their basic strengths as:

$$NH_3 < C_2H_5NH_2 < (C_2H_5)_5 NH$$

Again, $C_6H_5NH_2$ has proton acceptability less than NH_3 . Thus, we have:

$$C_6H_5NH_2 < NH_3 < C_2H_5NH_2 < (C_2H_5)_2 NH$$

Due to the – I effect of C_5H_5 group, the electron density on the N-atom in C_5H_5CH_2NH_2 is lower than that on the N-atom in C_2H_5NH_2 , but more than that in NH_3 . Therefore, the given compounds can karranged in the order of their basic strengths as:

$$C_6H_5NH_2 < NH_3 < C_6H_5CH_2NH_2 < C_2H_5NH_2 < (C_2H_5), NH$$

(ii) Considering the inductive effect and the steric hindrance of the alky groups. $C_2H_5NH_2$, $(C_2H_5)2NH_2$ and their basic strengths as follows $C_2H_1NH_2 < (C_2H_2)_2 N < (C_2H_2)_2 NH$

Again, due to the – R effect of C_0H_5 group, the electron density on the N atom in C_0H_5NH_2 is lower than that on the N atom in C_2H_5NH_2 . Therefore, the basicity of C_6H_5NH_2 is lower than that of ${}^{\rm C_2H_5NH_2}$. Hence, the given compounds can be arranged in the increasing order of their basic strengths as follows:

$${\rm C_6H_5NH_2} < {\rm C_2H_5NH_2} < \left({\rm \,C_2H_5} \right)_3 {\rm \,N} < \left({\rm \,C_2H_5} \right)_2 {\rm \,NH}$$

(iii) Considering the inductive effect and the steric hindrance of alkyl groups, $^{\text{CH}_3\text{NH}_2,(\text{CH}_3)_2\text{NH}}$ and $.^{(CH_3)_3}N$ can be arranged in the increasing order of their basic strengths as: (CH₃)₃ N < CH₃NH₂ < (CH₃), NH

In $C_6H_5NH_2$, N is directly attached to the benzene ring. Thus, the lone pair of electrons on the N – atom is delocalized over the benzene ring. In $C_6H_3CH_2NH_2$, N is not directly attached to the benzen ring. Thus, its lone pair is not delocalized over the benzene ring. Therefore, the electrons on the N atom are more easily available for protonation in C_6H_5CH_2NH_2 than in C_6H_5NH_2 i.e., C_6H_5CH_2NH_2 is more basic than C6H5NH3

Again, due to the – I effect of $C_5H_5CH_2NH_2$ is lower than that on the N vatom in $(CH_3)_3N$. Therefore, $(CH_3)_3N$ is more basic than $(C_6H_5CH_2NH_2)_3N$. Thus, the given compounds can be arranged in the increasing order of their basic strengths as follows.

$${\rm C_6H_5} < {\rm C_6H_5CH_2NH_2} < {\rm (CH_9)_3} \; {\rm N} < {\rm CH_3NH_2} < {\rm (CH_3)_2} \; {\rm NH}$$

Ans 3. The structures of different isomers corresponding to the molecular formula, C_3H_9N are given below:

(a)
$$CH_3 - CH_2 - CH_2 - NH_2$$

Propan-1-amine (1°)

(b)

$$\begin{array}{c} NH_2 \\ CH_3 - CH - CH_3 \\ \end{array}$$
 Propan-2-amine
$$\begin{array}{c} (1^\circ) \\ \text{(c)} \\ CH_3 - NH - C_2H_5 \\ N-\text{Methylethana} \min_{e \in 2^\circ)} \\ \text{(d)} \\ CH_3 \\ CH_3 - N - CH_3 \\ \end{array}$$

N, N-Dimethylmethanamine (3°)

^{1°} amines, (a) propan-1-amine, and (b) Propan-2-amine will liberate nitrogen gas on treatment with nitrous acid.

$$\begin{array}{ccc} \text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2 + \text{HNO}_2 & \longrightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + \text{N}_2 + \text{HCL} \\ \text{Propan}-1-\text{amine} & \text{Propan}-1-\text{ol} \\ \\ \text{CH}_3 - \text{CH} - \text{CH}_3 + \text{HNO}_2 & \longrightarrow \text{CH}_3 - \text{CH} - \text{CH}_3 + \text{N}_2 + \text{HCl} \\ \\ \text{NH}_2 & \text{OH} \\ \\ \text{Propan}-2-\text{amine} & \text{Propan}-2-\text{ol} \end{array}$$

Ans 4. (i) 1-Methylethanamine (1° amine)

- (ii) Propan-1-amine (1° amine)
- (iii) N-Methyl-2-methylethanamine (2° amine)
- (iv) 2-Methylpropan-2-amine (1° amine)
- (v) N-Methylbenzamine or N-methylaniline (2° amine)
- (vi) N-Ethyl-N-methylethanamine (3° amine)
- (vii) 3-Bromobenzenamine or 3-bromoaniline (1° amine)
- Ans 5. (i) Methylamine and dimethylamine can be distinguished by the carbylamine test. Carbylamine test: Aliphatic and aromatic primary amines on heating with chloroform and ethanolic potassium hydroxide form foul-smelling isocyanides or carbylamines. Methylamine (being an aliphatic primary amine) gives a positive carbylamine test, but dimethylamine does not.
- (ii) Secondary and tertiary amines can be distinguished by allowing them to react with Hinsberg's reagent (benzenesulphonyl chloride, C_6H_5SO_2Cl).

Secondary amines react with Hinsberg's reagent to form a product that is insoluble in an alkali. For example, N, N – diethylamine reacts with Hinsberg's reagent to form N, N – diethylbenzenesulphonamide, which is insoluble in an alkali. Tertiary amines, however, do not react with Hinsberg's reagent.

(iii) Ethylamine and aniline can be distinguished using the azo-dye test. A dye is obtained when aromatic amines react with ${}^{HNO_2\left(NaNO_2+dil.HCl\right)}$ at ${}^{0-5}{}^{\circ}C$, followed by a reaction with the alkalis solution of 2-naphthol. The dye is usually yellow, red, or orange in colour. Aliphatic amines give a brisk

effervescence due (to the evolution of N_2 gas) under similar conditions.

(iv) Aniline and benzylamine can be distinguished by their reactions with the help of nitrous acid, which is prepared in situ from a mineral acid and sodium nitrite. Benzylamine reacts with nitrous a to form unstable diazonium salt, which in turn gives alcohol with the evolution of nitrogen gas.

$$\begin{array}{c} C_6H_5CH_2-NH_2+HNO_2 & \underbrace{NaNO_2+HCI}_{C_6H_5CH_2}-N_2^{\dagger}C\overline{I} \\ Benzylamine & \underbrace{Uinstable}_{H_2O} \\ \\ N_2\uparrow+C_6H_5CH_2-OH+HCI \\ Benzyl alcobol & \\ \end{array}$$

On the other hand, aniline reacts with $^{\mathrm{HNO}_2}$ at a low temperature to form stable diazonium salt. Thus, nitrogen gas is not evolved.

$$C_gH_gNH_2\xrightarrow{NaNO_g+HCl}C_gH_g-N_2^+\bar{Cl}+NaCl+2H_3O$$

(v) Aniline and N-methylaniline can be distinguished using the Carbylamine test. Primary amines, o heating with chloroform and ethanolic potassium hydroxide, form foul-smelling isocyanides or carbylamines. Aniline, being an aromatic primary amine, gives positive carbylamine test. However, methylaniline, being a secondary amine does not.

Ans 6. (i) pKbof aniline is more than that of methylamine:

Aniline undergoes resonance and as a result, the electrons on the N-atom are delocalized over the benzene ring. Therefore, the electrons on the N-atom are less available to donate.

On the other hand, in case of methylamine (due to the +1 effect of methyl group), the electron density on the N-atom is increased. As a result, aniline is less basic than methylamine. Thus, pKb of aniline is more than that of methylamine.

(ii) Ethylamine is soluble in water whereas aniline is not:

Ethylamine when added to water forms intermolecular H – bonds with water. Hence, it is soluble it water.

SCIENCE

AMINES

But aniline does not undergo H – bonding with water to a very large extent due to the presence of large hydrophobic – C6H5 group. Hence, aniline is insoluble in water



(iii)Methylamine in water reacts with ferric chloride to precipitate hydrated ferric oxide:

Due to the +I effect of $-CH_3$ group, methylamine is more basic than water. Therefore, in water, methylamine produces OH – ions by accepting H+ ions from water.

$$CH_3 - NH_2 + H - OH \longrightarrow CH_3 - NH_3 + OH^-$$

Ferric chloride $^{(FeCl_3)}$ dissociates in water to form $^{Fe^-}$ and $^{CL^-}$ ions.

$$FeCl_3 \longrightarrow Fe^{3+} + 3Cl^{-}$$

Then, OH – ion reacts with Fe3+ ion to form a precipitate of hydrated ferric oxide.

(iv)Although amino group is o, p – directing in aromatic electrophilic substitution reactions, anilia on nitration gives a substantial amount of *m*-nitroaniline:

Nitration is carried out in an acidic medium. In an acidic medium, aniline is protonated to give anilinium ion (which is meta-directing).

For this reason, aniline on nitration gives a substantial amount of m-nitroaniline.


(v)Aniline does not undergo Friedel-Crafts reaction:

A Friedel-Crafts reaction is carried out in the presence of AlCl_3 . But AlCl_3 is acidic in nature, while aniline is a strong base. Thus, aniline reacts with AlCl₃ to form a salt (as shown in the following equation).

Due to the positive charge on the N-atom, electrophilic substitution in the benzene ring is deactivated. Hence, aniline does not undergo the Friedel-Crafts reaction.

(vi)Diazonium salts of aromatic amines are more stable than those of aliphatic amines:

The diazonium ion undergoes resonance as shown below:



This resonance accounts for the stability of the diazonium ion. Hence, diazonium salts of aromatic amines are more stable than those of aliphatic amines.

(vii)Gabriel phthalimide synthesis is preferred for synthesising primary amines:

Gabriel phthalimide synthesis results in the formation of amine only. 2 ° or 3 ° amines are not forme in this synthesis. Thus, a pure 1 ° amine can be obtained. Therefore, Gabriel phthalimide synthesis preferred for synthesizing primary amines.

Ans 7. (i) $\ln^{C_2H_5NH_2}$, only one $-^{C_2H_5}$ group is present while $\ln^{(C_2H_5)_2NH}$, two $-^{C_2H_5}$ groups are present. Thus, the +I effect is more in $(C_2H_5)_2NH$ than in $C_2H_5NH_2$. Therefore, the electron density over the N-atom is more in $(C_2H_5)_2NH$ than in $(C_2H_5NH_2)_2NH$ is more basic than $(C_2H_5NH_2)_2NH_2$.

Also, both C_6H_5NHCH_3 and ${}^{C_6H_5NH}_2$ are less basic than ${}^{(C_2H_5)_2NH}$ and ${}^{C_2H_5NH}_2$ due to the delocalization of the lone pair in the former two, Further, among C_6H_5NHCH_3 and ${}^{C_6H_5NH}_2$, the former will be more basic due to the +T effect of $-{}^{CH_3}$ group. Hence, the order of increasing basic of the given compounds is as follows:


$$C_6H_5NH_2 < C_6H_5NHCH_3 < C_2H_5NH_2 < C_2H_5$$
 NH

We know that the higher the basic strength, the lower is the pKb values.

$$C_6H_5NH_2 > C_6H_5NHCH_3 > C_2H_5NH_2 > (C_1H_5)_2NH$$

(ii) $C_6H_5N(CH_3)_2$ is more basic than $C_6H_5NH_2$ due to the presence of the +I effect of two -CH3 grou in $C_6H_5N(CH_3)_2$. Further, CH_3NH contains one $-CH_3$ group while C_2H_5 groups. Thus, C_2H_3NH is more basic than $C_2H_5NH_2$.

Now, $C_6H_5N(CH_3)_2$ is less basic than CH3NH2 because of the-R effect of $-C_6H_5$ group. Hence, the increasing order of the basic strengths of the given compounds is as follows: $C_6H_5NH_2 < C_6H_5NH_2 < C_6H_5NH_2 < C_6H_5NH_2 < C_6H_5NH_2 < C_7H_5 > 0$



SCIENCE

In p-toluidine, the presence of electron-donating $-^{CH_3}$ group increases the electron density on the N-atom.

Thus, p-toluidine is more basic than aniline.

On the other hand, the presence of electron-withdrawing

 $-^{NO_2}$ group decreases the electron density over the N-atom in p-nitroaniline. Thus, p-nitroaniline i less basic than aniline.

Hence, the increasing order of the basic strengths of the given compounds is as follows: p-Nitroaniline< Aniline < p-Toluidine

(b) $C_6H_5NHCH_3$ is more basic than $C_8H_5NH_2$ due to the presence of electron-donating $-CH_3$ group in $C_6H_5NHCH_3$

Again, in C_6H_5NHCH_3 , $-{}^{C_6H_5}$ group is directly attached to the N-atom. However, it is not so in C_6H_5NH_2NH_2 . Thus, in C_6H_5NHCH_3 , the -R effect of C_6H_5 group decreases the electron density over the N-atom. Therefore, C_6H_5NHCH_3 is more basic than C_6H_5NHCH_3 .

Hence, the increasing order of the basic strengths of the given compounds is as follows: $C_{\xi}H_{\xi}NH_{2} < C_{\xi}H_{\xi}NHCH_{3} < C_{\xi}H_{\xi}NH_{2}NH_{2}$

(iv) In the gas phase, there is no solvation effect. As a result, the basic strength mainly depends up the +I effect. The higher the +I effect, the stronger is the base. Also, the greater the number of alky groups, the higher is the +I effect. Therefore, the given compounds can be arranged in the decreasing order of their basic strengths in the gas phase as follows:


 $(C_2H_5)_3 N (C_2H_5)_3 NH C_2H_5 NH_2 NH_0$


(v) The boiling points of compounds depend on the extent of H-bonding present in that compound The more extensive the H-bonding in the compound, the higher is the boiling point. $(CH_3)_2 NH$ contains only one H-atom whereas $C_1H_5NH_2$ contains two H-atoms. Then, $C_2H_5NH_2$ undergoes more extensive H-bonding than $(CH_3)_2 NH$. Hence, the boiling point of $C_2H_5NH_2$ is higher than that of $(CH_3)_2 NH$.

Further, O is more electronegative than N. Thus, C_2H_5OH forms stronger H-bonds than C_2H_5NH_2 . A result, the boiling point of C_2H_5OH is higher than that of C_2H_5NH_2 and ${}^{(CH_3)_2}{}^{NH}$. Now, the given compounds can be arranged in the increasing order of their boiling points as follow $(CH_3)_2{}^{NH} < C_2H_5NH_2 < C_2H_5OH$

(vi) The more extensive the H-bonding, the higher is the solubility. C_2H_5NH_2 contains two H-atoms whereas ${}^{(C_2H_5)_2NH}$ contains only one H-atom. Thus, C_2H_5NH_2 undergoes more extensive H-bonding than ${}^{(C_2H_5)_2NH}$. Hence, the solubility in water of C_2H_5NH_2 is more than that of ${}^{(C_2H_5)_2NH}$. Further, the solubility of amines decreases with increase in the molecular mass. This is because the molecular mass of amines increases with an increase in the size of the hydrophobic part. The

molecular mass of C_6H_5NH_2 is greater than that of C_2H_5NH_2 and ${}^{(C_2H_5)_2NH}$. Hence, the increasing order of their solubility in water is as follows: ${}^{C_6H_5NH_2} < {}^{(C_2H_5)_2NH} < {}^{C_2H_5NH_2}$





Assertion and Reason Answers

1. (a) Assertion and reason both are correct statements and reason is correct explanation for assertion.


Explanation:

Ortho effect is a consequence of steric and electronic factors.

2. (d) Assertion is wrong statement but reason is correct statement.

Explanation:

In strongly acidic medium, aniline gets protonated and so the lone pair of electrons is not available to produce +E or + M effects. On the other hand, the -N+H3-N+H3 group exerts strong -I effect and thus it causes the deactivation of the ring.



Case Study Answers-

1. Answer:

- (i) (d) Azo dye test.
- (ii) (a) Aniline.

Explanation:

Aromatic primary amines give dye test.

(iii) (c) p-nitroaniline.

Explanation:

$$\begin{array}{c} \text{NH}_2 \\ \text{Aniline} \end{array} \xrightarrow{\text{CH}_3\text{COCl}} \xrightarrow{\text{NHCOCH}_3} \xrightarrow{\text{NHCOCH}_3} \\ \text{NH}_2 \\ \text{NO}_2 \\ \text{NO}_2 \\ p\text{-Nitroaniline} \end{array}$$

(c) p-benzoquinone. (iv)

Explanation:

$$\begin{array}{c}
NH_2 \\
\hline
MnO_2/H_2SO_4
\end{array}$$

$$\begin{array}{c}
O \\
O \\
O \\
P
-Benzoquinone$$

(v) (b) m-nitroaniline.

Explanation:

In acidic medium aniline gets protonated to anilinium ion which is meta-directing.

$$\begin{array}{c} \text{NH}_2 \\ \\ \text{NH}_2 \\ \\ \text{H}^+ \\ \\ \text{Aniline} \\ \\ \text{Anilinium ion} \\ \end{array} \begin{array}{c} \text{NH}_3 \\ \\ \text{NH}_3 \\ \\ \\ \text{NH}_3 \\ \\ \\ \text{NH}_2 \\ \\ \\ \text{NO}_2 \\ \\ \\ \\ \text{M-Nitroaniline} \\ \end{array}$$

2. Answer:

(i) (d) Dimethyl amine.

Explanation:

The increasing order of basicity of the given compounds is $(CH_3)_2NH > CH_3NH_2 > (CH_3)_3N > C_6H_5NH_2$. Due to the +I effect of alkyl groups, the electron density on nitrogen increases and thus, the availability of the lone pair of electrons to proton increases and hence, the basicity of amines also increases. So, aliphatic amines are more basic than aniline. In case of tertiary amine $(CH_3)_3N$, the covering of alkyl groups over nitrogen atom from all sides makes the approach and bonding by a proton relatively difficult, hence the basicity decreases. Electron withdrawing groups decrease electron density on nitrogen atom and thereby decreasing basicity.

(ii) (d) o-toluidine < aniline < m-toluidine

Explanation:

In general, electron donating (+R) group which when present on benzene ring (-NH₂, -OR, -R, etc.) at the para position increases the basicity of aniline.

Ortho substituted anilines are weaker bases than aniline due to ortho effect.

SCIENCE

AMINES

$$NH_2$$
 NH_2 NH_2 CH_3
 m -Toluidine Aniline o -Toluidine

(iii) (d) $(C_2H_5)_2NH > (C_2H_5)_3N > C_2H_5NH_2 > NH_3$

Explanation:

In case of ethylamines, the combined effect of (c) inductive effect, steric effect and salvation effect gives the order of basic strength as

$$({
m C}_2{
m H}_5)_2{
m NH} > ({
m C}_2{
m H}_5)_3{
m N} > {
m C}_2{
m H}_5{
m NH}_2 > {
m NH}_3 \ (2^\circ) \qquad (1^\circ)$$

(iv) (b) $(CH_3)_3N > (CH_3)_2NH > CH_3NH_2$

Explanation:

In non-aqueous solvents the basic strength of alkyl amines follows the order:

tertiary amines> secondary amines> primary amines.

(v) (c) Methylamine is a stronger base than ammonia.

Explanation:

Methyl amine is stronger base than ammonia due to electron releasing inductive effect of methyl group.