

Application of Derivatives

Question 1.

Find all the points of local maxima and local minima of the function $f(x) = (x - 1)^3 (x + 1)^2$

(a) 1, -1, -1/5

(b) 1, -1

(c) 1, -1/5

(d) -1, -1/5

Answer:

(a) 1, -1, -1/5

Question 2.

Find the local minimum value of the function $f(x) = \sin^4 x + \cos^4 x$, $0 < x < \frac{\pi}{2}$

(a) $\frac{1}{\sqrt{2}}$

(b) $\frac{1}{2}$

(c) $\frac{\sqrt{3}}{2}$

(d) 0

Answer:

(b) $\frac{1}{2}$

Question 3.

Find the points of local maxima and local minima respectively for the function $f(x) = \sin 2x - x$, where

$$-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$$

(a) $-\frac{\pi}{6}, \frac{\pi}{6}$

(b) $\frac{\pi}{3}, -\frac{\pi}{3}$

(c) $-\frac{\pi}{3}, \frac{\pi}{3}$

(d) $\frac{\pi}{6}, -\frac{\pi}{6}$

Answer:

(d) $\frac{\pi}{6}, -\frac{\pi}{6}$

Question 4.

If $y = \frac{ax-b}{(x-1)(x-4)}$ has a turning point P(2, -1), then find the value of a and b respectively.

- (a) 1, 2
- (b) 2, 1
- (c) 0, 1
- (d) 1, 0

Answer:

- (d) 1, 0

Question 5.

$\sin^p \theta \cos^q \theta$ attains a maximum, when $\theta =$

- (a) $\tan^{-1} \sqrt{\frac{p}{q}}$
- (b) $\tan^{-1} \left(\frac{p}{q} \right)$
- (c) $\tan^{-1} q$
- (d) $\tan^{-1} \left(\frac{q}{p} \right)$

Answer:

- (a) $\tan^{-1} \sqrt{\frac{p}{q}}$

Question 6.

Find the maximum profit that a company can make, if the profit function is given by $P(x) = 41 + 24x - 18x^2$.

- (a) 25
- (b) 43
- (c) 62
- (d) 49

Answer:

- (d) 49

Question 7.

If $y = x^3 + x^2 + x + 1$, then y

- (a) has a local minimum
- (b) has a local maximum
- (c) neither has a local minimum nor local maximum
- (d) None of these

Answer:

- (c) neither has a local minimum nor local maximum

Question 8.

Find both the maximum and minimum values respectively of $3x^4 - 8x^3 + 12x^2 - 48x + 1$ on the

interval $[1, 4]$.

- (a) -63, 257
- (b) 257, -40
- (c) 257, -63
- (d) 63, -257

Answer:

- (c) 257, -63

Question 9.

It is given that at $x = 1$, the function $x^4 - 62x^2 + ax + 9$ attains its maximum value on the interval $[0, 2]$. Find the value of a .

- (a) 100
- (b) 120
- (c) 140
- (d) 160

Answer:

- (b) 120

Question 10.

The function $f(x) = x^5 - 5x^4 + 5x^3 - 1$ has

- (a) one minima and two maxima
- (b) two minima and one maxima
- (c) two minima and two maxima
- (d) one minima and one maxima

Answer:

- (d) one minima and one maxima

Question 11.

Find the height of the cylinder of maximum volume that can be inscribed in a sphere of radius a .

- (a) $\frac{2a}{3}$
- (b) $\frac{2a}{\sqrt{3}}$
- (c) $\frac{a}{3}$
- (d) $\frac{a}{\sqrt{3}}$

Answer:

- (b) $\frac{2a}{\sqrt{3}}$

Question 12.

Find the volume of the largest cylinder that can be inscribed in a sphere of radius r cm.

- (a) $\frac{\pi r^3}{3\sqrt{3}}$
- (b) $\frac{4\pi r^2 h}{3\sqrt{3}}$

- (c) $4\pi r^3$
(d) $\frac{4\pi r^3}{3\sqrt{3}}$

Answer:

- (d) $\frac{4\pi r^3}{3\sqrt{3}}$

Question 13.

The area of a right-angled triangle of the given hypotenuse is maximum when the triangle is

- (a) scalene
(b) equilateral
(c) isosceles
(d) None of these

Answer:

- (c) isosceles

Question 14.

Find the area of the largest isosceles triangle having perimeter 18 metres.

- (a) $9\sqrt{3}$
(b) $8\sqrt{3}$
(c) $4\sqrt{3}$
(d) $7\sqrt{3}$

Answer:

- (a) $9\sqrt{3}$

Question 15.

$2x^3 - 6x + 5$ is an increasing function, if

- (a) $0 < x < 1$
(b) $-1 < x < 1$
(c) $x < -1$ or $x > 1$
(d) $-1 < x < -\frac{1}{2}$

Answer:

- (c) $x < -1$ or $x > 1$

Question 16.

If $f(x) = \sin x - \cos x$, then interval in which function is decreasing in $0 \leq x \leq 2\pi$, is

- (a) $\left[\frac{5\pi}{6}, \frac{3\pi}{4}\right]$
(b) $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$
(c) $\left[\frac{3\pi}{2}, \frac{5\pi}{2}\right]$

- (d) None of these

Answer:

(d) None of these

Question 17.

The function which is neither decreasing nor increasing in $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$ is

(a) cosec x

(b) tan x

(c) x^2

(d) $|x - 1|$

Answer:

(a) cosec x

Question 18.

The function $f(x) = \tan^{-1}(\sin x + \cos x)$ is an increasing function in

(a) $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

(b) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

(c) $\left(0, \frac{\pi}{2}\right)$

(d) None of these

Answer:

(d) None of these

Question 19.

The function $f(x) = x^3 + 6x^2 + (9 + 2k)x + 1$ is strictly increasing for all x, if

(a) $k > \frac{3}{2}$

(b) $k < \frac{3}{2}$

(c) $k \geq \frac{3}{2}$

(d) $k \leq \frac{3}{2}$

Answer:

(a) $k > \frac{3}{2}$

Question 20.

The point on the curves $y = (x - 3)^2$ where the tangent is parallel to the chord joining (3, 0) and (4, 1) is

(a) $\left(-\frac{7}{2}, \frac{1}{4}\right)$

(b) $\left(\frac{5}{2}, \frac{1}{4}\right)$

(c) $\left(-\frac{5}{2}, \frac{1}{4}\right)$

(d) $\left(\frac{7}{2}, \frac{1}{4}\right)$

Answer:

(d) $\left(\frac{7}{2}, \frac{1}{4}\right)$

Question 21.

The slope of the tangent to the curve $x = a \sin t$, $y = a\{\cot t + \log(\tan \frac{t}{2})\}$ at the point 't' is

- (a) $\tan t$
- (b) $\cot t$
- (c) $\tan \frac{t}{2}$
- (d) None of these

Answer:

- (a) $\tan t$

Question 22.

The equation of the normal to the curves $y = \sin x$ at $(0, 0)$ is

- (a) $x = 0$
- (b) $x + y = 0$
- (c) $y = 0$
- (d) $x - y = 0$

Answer:

- (b) $x + y = 0$

Question 23.

The tangent to the parabola $x^2 = 2y$ at the point $(1, \frac{1}{2})$ makes with the x-axis an angle of

- (a) 0°
- (b) 45°
- (c) 30°
- (d) 60°

Answer:

- (b) 45°

Question 24.

The two curves $x^3 - 3xy^2 + 5 = 0$ and $3x^2y - y^3 - 7 = 0$

- (a) cut at right angles
- (b) touch each other
- (c) cut at an angle $\frac{\pi}{4}$
- (d) cut at an angle $\frac{\pi}{3}$

Answer:

- (a) cut at right angles

Question 25.

The distance between the point (1, 1) and the tangent to the curve $y = e^{2x} + x^2$ drawn at the point $x = 0$

- (a) $\frac{1}{\sqrt{5}}$
- (b) $\frac{-1}{\sqrt{5}}$
- (c) $\frac{2}{\sqrt{5}}$
- (d) $\frac{-2}{\sqrt{5}}$

Answer:

- (c) $\frac{2}{\sqrt{5}}$

Question 26.

The tangent to the curve $y = 2x^2 - x + 1$ is parallel to the line $y = 3x + 9$ at the point

- (a) (2, 3)
- (b) (2, -1)
- (c) (2, 1)
- (d) (1, 2)

Answer:

- (d) (1, 2)

Question 27.

The tangent to the curve $y = x^2 + 3x$ will pass through the point (0, -9) if it is drawn at the point

- (a) (0, 1)
- (b) (-3, 0)
- (c) (-4, 4)
- (d) (1, 4)

Answer:

- (b) (-3, 0)

Question 28.

Find a point on the curve $y = (x - 2)^2$. at which the tangent is parallel to the chord joining the points (2, 0) and (4, 4).

- (a) (3, 1)
- (b) (4, 1)
- (c) (6, 1)
- (d) (5, 1)

Answer:

- (a) (3, 1)

Question 29.

Tangents to the curve $x^2 + y^2 = 2$ at the points (1, 1) and (-1, 1) are

- (a) parallel
- (b) perpendicular
- (c) intersecting but not at right angles
- (d) none of these

Answer:

- (b) perpendicular

Question 30.

If there is an error of 2% in measuring the length of a simple pendulum, then percentage error in its period is

- (a) 1%
- (b) 2%
- (c) 3%
- (d) 4%

Answer:

- (a) 1%

Question 31.

If there is an error of $a\%$ in measuring the edge of a cube, then percentage error in its surface area is

- (a) $2a\%$
- (b) $\frac{a}{2}\%$
- (c) $3a\%$
- (d) None of these

Answer:

- (b) $\frac{a}{2}\%$

Question 32.

If the radius of a sphere is measured as 9 cm with an error of 0.03 cm, then find the approximating error in calculating its volume.

- (a) $2.46\pi \text{ cm}^3$
- (b) $8.62\pi \text{ cm}^3$
- (c) $9.72\pi \text{ cm}^3$
- (d) $7.46\pi \text{ cm}^3$

Answer:

- (c) $9.72\pi \text{ cm}^3$

Question 33.

Find the approximate value of $f(3.02)$, where $f(x) = 3x^2 + 5x + 3$

- (a) 45.46
- (b) 45.76
- (c) 44.76

(d) 44.46

Answer:

(a) 45.46

Question 34.

$$f(x) = 3x^2 + 6x + 8, x \in \mathbb{R}$$

(a) 2

(b) 5

(c) -8

(d) does not exist

Answer:

(d) does not exist

Question 35.

The radius of a cylinder is increasing at the rate of 3 m/s and its height is decreasing at the rate of 4 m/s. The rate of change of volume when the radius is 4 m and height is 6 m, is

(a) 80π cu m/s

(b) 144π cu m/s

(c) 80 cu m/s

(d) 64 cu m/s

Answer:

(a) 80π cu m/s

Question 36.

The sides of an equilateral triangle are increasing at the rate of 2 cm/s. The rate at which the area increases, when the side is 10 cm, is

(a) $\sqrt{3}$ cm²/s

(b) 10 cm²/s

(c) $10\sqrt{3}$ cm²/s

(d) $\frac{10}{\sqrt{3}}$ cm²/s

Answer:

(c) $10\sqrt{3}$ cm²/s

Question 37.

A particle is moving along the curve $x = at^2 + bt + c$. If $ac = b^2$, then particle would be moving with uniform

(a) rotation

(b) velocity

(c) acceleration

(d) retardation

Answer:

(c) acceleration

Question 38.

The distance 's' metres covered by a body in t seconds, is given by $s = 3t^2 - 8t + 5$. The body will stop after

(a) 1 s

(b) $\frac{3}{4}$ s

(c) $\frac{4}{3}$ s

(d) 4 s

Answer:

(c) $\frac{4}{3}$ s

Question 39.

The position of a point in time 't' is given by $x = a + bt - ct^2$, $y = at + bt^2$. Its acceleration at time 't' is

(a) $b - c$

(b) $b + c$

(c) $2b - 2c$

(d) $2\sqrt{b^2 + c^2}$

Answer:

(d) $2\sqrt{b^2 + c^2}$

Question 40.

The function $f(x) = \log(1 + x) - \frac{2x}{2+x}$ is increasing on

(a) $(-1, \infty)$

(b) $(-\infty, 0)$

(c) $(-\infty, \infty)$

(d) None of these

Answer:

(a) $(-1, \infty)$

Question 41.

$f(x) = \left(\frac{e^{2x}-1}{e^{2x}+1} \right)$ is

(a) an increasing function

(b) a decreasing function

(c) an even function

(d) None of these

Answer:

(a) an increasing function

Question 42.

The function $f(x) = \cot^{-1} x + x$ increases in the interval

- (a) $(1, \infty)$
- (b) $(-1, \infty)$
- (c) $(0, \infty)$
- (d) $(-\infty, \infty)$

Answer:

- (d) $(-\infty, \infty)$

Question 43.

The function $f(x) = \frac{x}{\log x}$ increases on the interval

- (a) $(0, \infty)$
- (b) $(0, e)$
- (c) (e, ∞)
- (d) none of these

Answer:

- (c) (e, ∞)

Question 44.

The length of the longest interval, in which the function $3 \sin x - 4 \sin^3 x$ is increasing, is

- (a) $\frac{\pi}{3}$
- (b) $\frac{\pi}{2}$
- (c) $\frac{3\pi}{2}$
- (d) π

Answer:

- (a) $\frac{\pi}{3}$

Question 45.

The coordinates of the point on the parabola $y^2 = 8x$ which is at minimum distance from the circle $x^2 + (y + 6)^2 = 1$ are

- (a) $(2, -4)$
- (b) $(18, -12)$
- (c) $(2, 4)$
- (d) none of these

Answer:

- (a) $(2, -4)$

Question 46.

The distance of that point on $y = x^4 + 3x^2 + 2x$ which is nearest to the line $y = 2x - 1$ is

- (a) $\frac{3}{\sqrt{5}}$

(b) $\frac{4}{\sqrt{5}}$

(c) $\frac{2}{\sqrt{5}}$

(d) $\frac{1}{\sqrt{5}}$

Answer:

(d) $\frac{1}{\sqrt{5}}$

Question 47.

The function $f(x) = x + \frac{4}{x}$ has

(a) a local maxima at $x = 2$ and local minima at $x = -2$

(b) local minima at $x = 2$, and local maxima at $x = -2$

(c) absolute maxima at $x = 2$ and absolute minima at $x = -2$

(d) absolute minima at $x = 2$ and absolute maxima at $x = -2$

Answer:

(b) local minima at $x = 2$, and local maxima at $x = -2$

Question 48.

The combined resistance R of two resistors R_1 and R_2 ($R_1, R_2 > 0$) is given by $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$. If

$R_1 + R_2 = C$ (a constant), then maximum resistance R is obtained if

(a) $R_1 > R_2$

(b) $R_1 < R_2$

(c) $R_1 = R_2$

(d) None of these

Answer:

(c) $R_1 = R_2$

Question 49.

Find the height of a cylinder, which is open at the top, having a given surface area, greatest volume and of radius r .

(a) r

(b) $2r$

(c) $\frac{r}{2}$

(d) $\frac{3\pi r}{2}$

Answer:

(a) r