


Unit 6 (Triangles)

Exercise 6.1 Multiple Choice Questions (MCQs)

Question 1:


In figure, if $\angle BAC = 90^\circ$ and $AD \perp BC$. Then,



- (a) $BD \cdot CD = BC^2$ (b) $AB \cdot AC = BC^2$ (c) $BD \cdot CD = AD^2$
 (d) $AB \cdot AC = AD^2$

Solution:

(c) In $\triangle ADB$ and $\triangle ADC$,



$\therefore$

$\therefore$

$\Rightarrow$

$$\begin{aligned}\angle D &= \angle D = 90^\circ \\ \angle DBA &= \angle DAC \\ \triangle ADB &\sim \triangle ADC \\ \frac{BD}{AD} &= \frac{AD}{CD} \\ BD \cdot CD &= AD^2\end{aligned}$$

[each equal to $90^\circ - \angle C$]
 [by AAA similarity criterion]

Question 2:

If the lengths of the diagonals of rhombus are 16 cm and 12 cm. Then, the length of the sides of the rhombus is

- (a) 9 cm (b) 10 cm (c) 8 cm (d) 20 cm

Solution:


(b) We know that, the diagonals of a rhombus are perpendicular bisector of each other.

Given, $AC = 16$ cm and $BD = 12$ cm [let]

$\therefore$ $AO = 8$ cm, $SO = 6$ cm

and $\angle AOB = 90^\circ$

In right angled $\triangle AOB$,



$$AB^2 = AO^2 + OB^2 \quad [\text{by Pythagoras theorem}]$$

$\Rightarrow$

$$AB^2 = 8^2 + 6^2 = 64 + 36 = 100$$

$\therefore$

$$AB = 10\text{cm}$$

Question 3:

If $\triangle ABC \sim \triangle EDF$ and $\triangle ABC$ is not similar to $\triangle DEF$, then which of the following is not true?


- (a) $BC \cdot EF = AC \cdot FD$ (b) $AB \cdot EF = AC \cdot DE$
 (c) $BC \cdot DE = AB \cdot EF$ (d) $BC \cdot DE = AB \cdot FD$

Solution:

(c) Given,

$$\triangle ABC \sim \triangle EDF$$

$$\frac{AB}{ED} = \frac{BC}{DF} = \frac{AC}{EF}$$



Taking first two terms, we get

$$\frac{AB}{ED} = \frac{BC}{DF}$$

$\Rightarrow$

$$AB \cdot DF = ED \cdot BC$$

or

$$BC \cdot DE = AB \cdot DF$$

So, option (d) is true.

Taking last two terms, we get

$$\frac{BC}{DF} = \frac{AC}{EF}$$

$\Rightarrow$

$$BC \cdot EF = AC \cdot DF$$

So, option (a) is also true.

Taking first and last terms, we get

$$\frac{AB}{ED} = \frac{AC}{EF}$$

$\Rightarrow$

$$AB \cdot EF = ED \cdot AC$$

Hence, option (b) is true.

Question 4:

If in two $\triangle PQR$, $\frac{AB}{QR} = \frac{BC}{PR} = \frac{CA}{PQ}$, then

- (a) $\triangle PQR \sim \triangle CAB$ (b) $\triangle PQR \sim \triangle ABC$
 (c) $\triangle CBA \sim \triangle PQR$ (d) $\triangle BCA \sim \triangle PQR$


Solution:

(a) Given, in two $\triangle ABC$ and $\triangle PQR$, $\frac{AB}{QR} = \frac{BC}{PR} = \frac{CA}{PQ}$

which shows that sides of one triangle are proportional to the side of the other triangle, then their corresponding angles are also equal, so by SSS similarity, triangles are similar.

i.e.,

$$\triangle CAB \sim \triangle PQR$$



Question 5:

In figure, two line segments AC and BD intersect each other at the point P such that PA = 6 cm, PB = 3 cm, PC = 2.5 cm, PD = 5 cm, $\angle APB = 50^\circ$ and $\angle CDP = 30^\circ$. Then, $\angle PBA$ is equal to

- (a) 50° (b) 30° (c) 60° (d) 100°

Solution:

(d) In $\triangle APB$ and $\triangle CPD$, $\angle APB = \angle CPD = 50^\circ$ [vertically opposite angles]
 $\frac{AP}{PD} = \frac{6}{5}$... (i)
 and $\frac{BP}{CP} = \frac{3}{2.5} = \frac{6}{5}$... (ii)
 From Eqs. (i) and (ii)
 $\frac{AP}{PD} = \frac{BP}{CP}$
 $\therefore \triangle APB \sim \triangle DPC$ [by SAS similarity criterion]
 $\therefore \angle A = \angle D = 30^\circ$ [corresponding angles of similar triangles]
 In $\triangle APB$, $\angle A + \angle B + \angle APB = 180^\circ$ [sum of angles of a triangle = 180°]
 $\Rightarrow 30^\circ + \angle B + 50^\circ = 180^\circ$
 $\therefore \angle B = 180^\circ - (50^\circ + 30^\circ) = 100^\circ$
 i.e., $\angle PBA = 100^\circ$


Question 6:

If in two $\triangle DEF$ and $\triangle PQR$, $\angle D = \angle Q$ and $\angle R = \angle E$, then which of the following is not true?

- (a) $\frac{EF}{PR} = \frac{DF}{PQ}$ (b) $\frac{DE}{PQ} = \frac{EF}{RP}$ (c) $\frac{DE}{QR} = \frac{DF}{PQ}$ (d) $\frac{EF}{RP} = \frac{DE}{QR}$

Solution:

(b) Given, in $\triangle DEF$, $\angle D = \angle Q$, $\angle R = \angle E$



$\therefore \triangle DEF \sim \triangle QRP$ [by AAA similarity criterion]
 $\Rightarrow \angle F = \angle P$ [corresponding angles of similar triangles]
 $\therefore \frac{DF}{QP} = \frac{ED}{RQ} = \frac{FE}{PR}$

Question 7:

In $\triangle ABC$ and $\triangle DEF$, $\angle B = \angle E$, $\angle F = \angle C$ and $AB = 30E$. Then, the two triangles are

- (a) congruent but not similar (b) similar but not congruent
 (c) neither congruent nor similar (d) congruent as well as similar

Solution:

Question 8:


Solution:

We know that, the ratio of the areas of two similar triangles is equal to square of the ratio of their corresponding sides.

Question 9:

(a) $DE = 12$ cm, $\angle F = 50^\circ$ (b) $DE = 12$ cm, $\angle F = 100^\circ$
(c) $EF = 12$ cm, $\angle D = 100^\circ$ (d) $EF = 12$ cm, $\angle D = 30^\circ$

(b) Given, $\triangle ABC \sim \triangle ADF$, then $\angle A = \angle D = 30^\circ$, $\angle C = \angle E = 50^\circ$




Question 10:

(a) $\angle B = \angle E$ (b) $\angle A = \angle D$
(c) $\angle B = \angle D$ (d) $\angle A = \angle F$

Solution:

(c) Given, in $\triangle ABC$ and $\triangle EDF$,

$$\frac{AB}{DE} = \frac{BC}{FD}$$



By converse of basic proportionality theorem,

$$\triangle ABC \sim \triangle EDF$$

Then,

$$\angle B = \angle D, \angle A = \angle E$$

and

$$\angle C = \angle F$$

Question 11:

If $\triangle ABC \sim \triangle QRP$, $\frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{9}{4}$, $AB = 18$ cm and $BC = 15$ cm, then PR is equal to

(a) 10 cm


(b) 12 cm

(c) $\frac{20}{3}$ cm

(d) 8 cm

Solution:

(a) Given, $\triangle ABC \sim \triangle QRP$, $AB = 18$ cm and $BC = 15$ cm



We know that, the ratio of area of two similar triangles is equal to the ratio of square of their corresponding sides.

$$\begin{aligned} \therefore \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle QRP)} &= \frac{(BC)^2}{(RP)^2} \\ \text{But given, } \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} &= \frac{9}{4} && \text{[given]} \\ \Rightarrow \frac{(15)^2}{(RP)^2} &= \frac{9}{4} && [\because BC = 15 \text{ cm, given}] \\ \Rightarrow (RP)^2 &= \frac{225 \times 4}{9} = 100 \\ \therefore RP &= 10 \text{ cm} \end{aligned}$$

Question 12:

If S is a point on side PQ of a $\triangle PQR$ such that $PS = QS = RS$, then

(a) $PR \cdot QR = RS^2$

(b) $QS^2 + RS^2 = QR^2$

(c) $PR^2 + QR^2 = PQ^2$

(d) $PS^2 + RS^2 = PR^2$

Solution:

(c) Given, in ΔPQR ,

$$PS = QS = RS \quad \dots (i)$$

$$\text{In } \Delta PSR, \quad PS = RS \quad [\text{from Eq. (i)}]$$

$$\Rightarrow \quad \angle 1 = \angle 2 \quad \dots (ii)$$

$$\text{Similarly, in } \Delta RSQ, \quad \dots$$

$$\Rightarrow \quad \angle 3 = \angle 4 \quad \dots (iii)$$

[corresponding angles of equal sides are equal]

Now, in ΔPQR , sum of angles = 180°

$$\Rightarrow \quad \angle P + \angle Q + \angle R = 180^\circ$$

$$\Rightarrow \quad \angle 2 + \angle 4 + \angle 1 + \angle 3 = 180^\circ$$

$$\Rightarrow \quad \angle 1 + \angle 3 + \angle 1 + \angle 3 = 180^\circ$$


$$\Rightarrow \quad 2(\angle 1 + \angle 3) = 180^\circ$$

$$\Rightarrow \quad \angle 1 + \angle 3 = \frac{180^\circ}{2} = 90^\circ$$

$$\therefore \quad \angle R = 90^\circ$$

In ΔPQR , by Pythagoras theorem,

$$PR^2 + QR^2 = PQ^2$$



[using Eqs. (ii) and (iii)]

Exercise 6.2 Very Short Answer Type Questions

Question 1:

Is the triangle with sides 25 cm, 5 cm and 24 cm a right triangle? Give reason for your answer.

Solution:

False

Let $a = 25$ cm, $b = 5$ cm and $c = 24$ cm

$$\begin{aligned} \text{Now,} \quad b^2 + c^2 &= (5)^2 + (24)^2 \\ &= 25 + 576 = 601 \neq (25)^2 \end{aligned}$$

Hence, given sides do not make a right triangle because it does not satisfy the property of Pythagoras theorem.

Question 2:

It is given that $\Delta DEF \sim \Delta RPQ$. Is it true to say that $\angle D = \angle R$ and $\angle F = \angle P$? Why?

Solution:

False

We know that, if two triangles are similar, then their corresponding angles are equal.

$$\therefore \quad \angle D = \angle R, \angle E = \angle P \text{ and } \angle F = \angle Q$$


Question 3:

A and B are respectively the points on the sides PQ and PR of a ΔPQR such that $PQ = 12.5$ cm, $PA = 5$ cm, $BR = 6$ cm and $PB = 4$ cm. Is $AB \parallel QR$? Give reason for your answer.

Solution:

False

Given, $PQ = 12.5$ cm, $PA = 5$ cm, $BR = 6$ cm and $PB = 4$ cm



Then,

$$QA = QP - PA = 12.5 - 5 = 7.5 \text{ cm}$$

Now,

$$\frac{PA}{AQ} = \frac{5}{7.5} = \frac{50}{75} = \frac{2}{3} \quad \dots(i)$$

and

$$\frac{PB}{BR} = \frac{4}{6} = \frac{2}{3} \quad \dots(ii)$$

From Eqs. (i) and (ii),


$$\frac{PA}{AQ} = \frac{PB}{BR}$$

By converse of basic proportionality theorem,

$$AB \parallel QR$$

Question 4:

In figure, BD and CE intersect each other at the point P. Is $\triangle PBC \sim \triangle PDE$? Why?



Solution:

True

$$\angle BPC = \angle EPD \quad [\text{vertically opposite angles}]$$

Now,

$$\frac{PB}{PD} = \frac{5}{10} = \frac{1}{2} \quad \dots(i)$$

and

$$\frac{PC}{PE} = \frac{6}{12} = \frac{1}{2} \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$\frac{PB}{PD} = \frac{PC}{PE}$$

Since, one angle of $\triangle PBC$ is equal to one angle of $\triangle PDE$ and the sides including these angles are proportional, so both triangles are similar.

Hence, $\triangle PBC \sim \triangle PDE$, by SAS similarity criterion.

Question 5:

In $\triangle PQR$ and $\triangle MST$, $\angle P = 55^\circ$, $\angle Q = 25^\circ$, $\angle M = 100^\circ$ and $\angle S = 25^\circ$. Is $\triangle PQR \sim \triangle TSM$? Why?

Solution:

False

We know that, the sum of three angles of a triangle is 180° .

In $\triangle PQR$,

$$\angle P + \angle Q + \angle R = 180^\circ$$

$\Rightarrow$

$$55^\circ + 25^\circ + \angle R = 180^\circ$$

$\Rightarrow$

$$\angle R = 180^\circ - (55^\circ + 25^\circ) = 180^\circ - 80^\circ = 100^\circ$$

In $\triangle TSM$,

$$\angle T + \angle S + \angle M = 180^\circ$$


$\Rightarrow$

$$\angle T + 25^\circ + 100^\circ = 180^\circ$$

$\Rightarrow$

$$\angle T = 180^\circ - (25^\circ + 100^\circ)$$

$$= 180^\circ - 125^\circ = 55^\circ$$



In ΔPQR and ΔTSM , and

$\angle P = \angle T$, $\angle Q = \angle S$,

and

$\angle R = \angle M$

$\Delta PQR \sim \Delta TSM$ [since, all corresponding angles are equal]

Hence, ΔPQR is not similar to ΔTSM , since correct correspondence is $P \leftrightarrow T$, $Q \leftrightarrow S$ and $R \leftrightarrow M$

Question 6:

Is the following statement true? Why? "Two quadrilaterals are similar, if their corresponding angles are equal".

Solution:

False

Two quadrilaterals are similar, if their corresponding angles are equal and corresponding sides must also be proportional.

Question 7:

Two sides and the perimeter of one triangle are respectively three times the corresponding sides and the perimeter of the other triangle. Are the two triangles similar? Why?

Solution:

True

Here, the corresponding two sides and the perimeters of two triangles are proportional, then third side of both triangles will also be in proportion.

Question 8:

If in two right triangles, one of the acute angles of one triangle is equal to an acute angle of the other triangle. Can you say that two triangles will be similar? Why?

Solution:

True

Let two right angled triangles be ΔABC and ΔPQR .

Question 9:

The ratio of the corresponding altitudes of two similar triangles is $\frac{3}{5}$. Is it correct to say that ratio of their areas is $\frac{6}{5}$? Why?

Solution:

False

By the property of area of two similar triangles,

$$\begin{aligned} \Rightarrow \left(\frac{\text{Area}_1}{\text{Area}_2} \right) &= \left(\frac{\text{Altitude}_1}{\text{Altitude}_2} \right)^2 \\ \left(\frac{\text{Area}_1}{\text{Area}_2} \right) &= \left(\frac{3}{5} \right)^2 \\ &= \frac{9}{25} \neq \frac{6}{5} \end{aligned} \quad \left[\because \frac{\text{altitude}_1}{\text{altitude}_2} = \frac{3}{5}, \text{ given} \right]$$

So, given statement is not correct,

Question 10:

D is a point on side QR of $\triangle PQR$ such that $PD \perp QR$. Will it be correct to say that $\triangle PQD \sim \triangle RPD$? Why?

Solution:

False


In $\triangle PQD$ and $\triangle RPD$,

$$PD = PD$$

[common side]

$$\angle PDQ = \angle PDR$$

[each 90°]



Here, no other sides or angles are equal, so we can say that $\triangle PQD$ is not similar to $\triangle RPD$.


But, if $\angle P = 90^\circ$,

then $\angle DPQ = \angle PRD$

[each equal to $90^\circ - \angle Q$ and by ASA similarity criterion, $\triangle PQD \sim \triangle RPD$]

Question 11:

In figure, if $\angle D = \angle C$, then it is true that $\triangle ADE \sim \triangle ACB$? Why?



Solution:

True

In $\triangle ADE$ and $\triangle ACB$,

$$\angle A = \angle A$$

[common angle]

$$\angle D = \angle C$$

[given]

$$\triangle ADE \sim \triangle ACB$$

[by AAA similarity criterion]

Question 12:

Is it true to say that, if in two triangles, an angle of one triangle is equal to an angle of another triangle and two sides of one triangle are proportional to the two sides of the other triangle, then the triangles are similar? Give reason for your answer.

Solution:

False

Because, according to SAS similarity criterion, if one angle of a triangle is equal to one angle of the other triangle and the sides including these angles are proportional, then the two triangles are similar.

Here, one angle and two sides of two triangles are equal but these sides not including equal angle, so given statement is not correct.

Exercise 6.3 Short Answer Type Questions

Question 1:

In a $\triangle PQR$, $PR^2 - PQ^2 = QR^2$ and M is a point on side PR such that $QM \perp PR$.

Prove that $QM^2 = PM \times MR$.

Solution:

Given In ΔPQR ,

$$PR^2 - PQ^2 = QR^2 \text{ and } QM \perp PR$$

To prove


$$QM^2 = PM \times MR$$

Proof Since,

$$PR^2 - PQ^2 = QR^2$$

$\Rightarrow$

$$PR^2 = PQ^2 + QR^2$$



So, ΔPQR is right angled triangle at Q.

In ΔQMR and ΔPMQ ,

$$\angle M = \angle M$$

[each 90°]

$$\angle MQR = \angle QPM$$

[each equal to $90^\circ - \angle R$]

$$\Delta QMR \sim \Delta PMQ$$

[by AAA similarity criterion]

Now, using property of area of similar triangles, we get

$$\frac{\text{ar}(\Delta QMR)}{\text{ar}(\Delta PMQ)} = \frac{(QM)^2}{(PM)^2}$$

$$\Rightarrow \frac{\frac{1}{2} \times RM \times QM}{\frac{1}{2} \times PM \times QM} = \frac{(QM)^2}{(PM)^2} \quad [\because \text{area of triangle} = \frac{1}{2} \times \text{base} \times \text{height}]$$

$$\Rightarrow QM^2 = PM \times RM \quad \text{Hence proved.}$$

Question 2:

Find the value of x for which $DE \parallel AB$ in given figure.

Solution:

Given,

$$\frac{DE \parallel AB}{\frac{CD}{AD} = \frac{CE}{BE}}$$

[by basic proportionality theorem]

$\Rightarrow$

$$\frac{x+3}{3x+19} = \frac{x}{3x+4}$$

$\Rightarrow$

$$(x+3)(3x+4) = x(3x+19)$$

$\Rightarrow$

$$3x^2 + 4x + 9x + 12 = 3x^2 + 19x$$

$\Rightarrow$


$$19x - 13x = 12$$

$\Rightarrow$

$$6x = 12$$

$\therefore$


$$x = \frac{12}{6} = 2$$



Hence, the required value of x is 2.

Question 3:

In figure, if $\angle 1 = \angle 2$ and $\Delta NSQ = \Delta MTR$, then prove that $\Delta PTS \sim \Delta PRQ$.



Solution:

Given $\triangle NSQ \cong \triangle MTR$ and $\angle 1 = \angle 2$

To prove $\triangle PTS \sim \triangle PRQ$

Proof Since,

$$\triangle NSQ \cong \triangle MTR$$

So,

$$SQ = TR \quad \dots (i)$$

Also,

$$\angle 1 = \angle 2 \Rightarrow PT = PS \quad \dots (ii)$$

[since, sides opposite to equal angles are also equal]

From Eqs. (i) and (ii),

$$\frac{PS}{SQ} = \frac{PT}{TR}$$

$\Rightarrow$

$$ST \parallel QR \quad [\text{by converse of basic proportionality theorem}]$$

$\therefore$

$$\angle 1 = \angle PQR$$

and

$$\angle 2 = \angle PRQ$$

In $\triangle PTS$ and $\triangle PRQ$,

[common angles]

$$\angle P = \angle P$$

$$\angle 1 = \angle PQR$$

$$\angle 2 = \angle PRQ$$

$\therefore$

$$\triangle PTS \sim \triangle PRQ$$

[by AAA similarity criterion]

Hence proved.

Question 4:

Diagonals of a trapezium PQRS intersect each other at the point O, $PQ \parallel RS$ and $PQ = 3$

RS. Find the ratio of the areas of


$\triangle POQ$ and $\triangle ROS$.

Solution:

Given PQRS is a trapezium in which $PQ \parallel RS$ and $PQ = 3$ RS

$\Rightarrow$

$$\frac{PQ}{RS} = \frac{3}{1} \quad \dots (i)$$



In $\triangle POQ$ and $\triangle ROS$,

$$\angle SOR = \angle QOP$$

[vertically opposite angles]

$$\angle SRP = \angle RPQ$$

[alternate angles]

$\therefore$

$$\triangle POQ \sim \triangle ROS$$

[by AAA similarity criterion]

By property of area of similar triangle,

$$\frac{\text{ar}(\triangle POQ)}{\text{ar}(\triangle ROS)} = \frac{(PQ)^2}{(RS)^2} = \left(\frac{PQ}{RS}\right)^2 = \left(\frac{3}{1}\right)^2 \quad [\text{from Eq. (i)}]$$


$\Rightarrow$

$$\frac{\text{ar}(\triangle POQ)}{\text{ar}(\triangle ROS)} = \frac{9}{1}$$

Hence, the required ratio is 9 : 1.

Question 5:

In figure, if $AB \parallel DC$ and AC, PQ intersect each other at the point O. Prove that $OA \cdot CQ = OC \cdot AP$.



Solution:

Given AC and PQ intersect each other at the point O and $AB \parallel DC$

Prove that $OA \cdot CQ = OC \cdot AP$.

Proof In $\triangle AOP$ and $\triangle COQ$, $\angle AOP = \angle COQ$ [vertically opposite angles]
 $\angle APO = \angle CQO$
[since, $AB \parallel DC$ and PQ is transversal, so alternate angles]
 $\therefore \triangle AOP \sim \triangle COQ$ [by AAA similarity criterion]
Then, $\frac{OA}{OC} = \frac{AP}{CQ}$ [since, corresponding sides are proportional]
 $\Rightarrow OA \cdot CQ = OC \cdot AP$ **Hence proved.**

Question 6:

Find the altitude of an equilateral triangle of side 8 cm.

Solution:

Let ABC be an equilateral triangle of side 8 cm i.e., $AB = BC = CA = 8$ cm. Draw altitude AD which is perpendicular to BC. Then, D is the mid-point of BC.

$$\therefore BD = CD = \frac{1}{2} BC = \frac{8}{2} = 4 \text{ cm}$$

$$\text{Now, } AB^2 = AD^2 + BD^2 \quad [\text{by Pythagoras theorem}]$$


$$\Rightarrow (8)^2 = AD^2 + (4)^2$$

$$\Rightarrow 64 = AD^2 + 16$$

$$\Rightarrow AD^2 = 64 - 16 = 48$$

$$\Rightarrow AD = \sqrt{48} = 4\sqrt{3} \text{ cm.}$$

Hence, altitude of an equilateral triangle is $4\sqrt{3}$ cm.



Question 7:

If $\triangle ABC \sim \triangle DEF$, $AB = 4$ cm, $DE = 6$, $EF = 9$ cm and $FD = 12$ cm, then find the perimeter of $\triangle ABC$.

Solution:

Given $AB = 4$ cm, $DE = 6$ cm and $EF = 9$ cm and $FD = 12$ cm

Also,

$$\begin{aligned} \therefore \triangle ABC &\sim \triangle DEF \\ \frac{AB}{ED} &= \frac{BC}{EF} = \frac{AC}{DF} \\ \Rightarrow \frac{4}{6} &= \frac{BC}{9} = \frac{AC}{12} \end{aligned}$$


On taking first two terms, we get

$$\begin{aligned} \Rightarrow \frac{4}{6} &= \frac{BC}{9} \\ BC &= \frac{4 \times 9}{6} = 6 \text{ cm} \\ &= AC = \frac{6 \times 12}{9} = 8 \text{ cm} \end{aligned}$$

$$\begin{aligned} \text{Now, perimeter of } \triangle ABC &= AB + BC + AC \\ &= 4 + 6 + 8 = 18 \text{ cm} \end{aligned}$$

Question 8:

In figure, if $DE \parallel BC$, then find the ratio of ar ($\triangle ADE$) and ar (DECB).



Solution:

Given, $DE \parallel BC$, $DE = 6$ cm and $BC = 12$ cm
In ΔABC and ΔADE ,

$$\angle ABC = \angle ADE$$

[corresponding angle]

$$\angle ACB = \angle AED$$

[corresponding angle]

and

$$\angle A = \angle A$$

[common side]

$\therefore$

$$\Delta ABC \sim \Delta AED$$

[by AAA similarity criterion]

Then,

$$\frac{\text{ar}(\Delta ADE)}{\text{ar}(\Delta ABC)} = \frac{(DE)^2}{(BC)^2}$$

$$= \frac{(6)^2}{(12)^2} = \left(\frac{1}{2}\right)^2$$

$\Rightarrow$

$$\frac{\text{ar}(\Delta ADE)}{\text{ar}(\Delta ABC)} = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$$

Let $\text{ar}(\Delta ADE) = k$, then $\text{ar}(\Delta ABC) = 4k$

Now, $\text{ar}(DECB) = \text{ar}(ABC) - \text{ar}(ADE) = 4k - k = 3k$


$\therefore$ Required ratio = $\text{ar}(ADE) : \text{ar}(DECB) = k : 3k = 1 : 3$

Question 9:

ABCD is a trapezium in which $AB \parallel DC$ and P, Q are points on AD and BC respectively, such that $PQ \parallel DC$, if $PD = 18$ cm, $BQ = 35$ cm and $QC = 15$ cm, find AD.

Solution:

Given, a trapezium ABCD in which $AB \parallel DC$. P and Q are points on AD and BC, respectively such that $PQ \parallel DC$. Thus, $AB \parallel PQ \parallel DC$.



Join BD .

In ΔABD ,

By basic proportionality theorem,

$$\frac{PO \parallel AB}{DP} = \frac{DO}{AP}$$

[$\because PQ \parallel AB$]

...(i)

In ΔBDC ,

By basic proportionality theorem,

$$OQ \parallel DC$$

[$\because PQ \parallel DC$]

$$\frac{BQ}{QC} = \frac{OB}{OD}$$

$$\frac{BQ}{QC} = \frac{OD}{OB}$$

...(ii)

$\Rightarrow$

From Eqs. (i) and (ii),

$$\frac{DP}{AP} = \frac{QC}{BQ}$$

$\Rightarrow$

$$\frac{18}{AP} = \frac{15}{35}$$

$\Rightarrow$

$$AP = \frac{18 \times 35}{15} = 42$$

$\therefore$

$$AD = AP + DP = 42 + 18 = 60 \text{ cm}$$

Question 10:

Corresponding sides of two similar triangles are in the ratio of 2 : 3. If the area of the smaller triangle is 48 cm^2 , then find the area of the larger triangle.

Solution:

Given, ratio of corresponding sides of two similar triangles = $2:3$ or $\frac{2}{3}$

Area of smaller triangle = 48 cm^2

By the property of area of two similar triangle,

Ratio of area of both triangles = (Ratio of their corresponding sides)²

$$\begin{aligned}
 \text{i.e.,} \quad & \frac{\text{ar (smaller triangle)}}{\text{ar (larger triangle)}} = \left(\frac{2}{3}\right)^2 \\
 \Rightarrow & \frac{48}{\text{ar(larger triangle)}} = \frac{4}{9} \\
 \Rightarrow & \text{ar (larger triangle)} = \frac{48 \times 9}{4} = 12 \times 9 = 108 \text{ cm}^2
 \end{aligned}$$

Question 11:

In a ΔPQR , N is a point on PR , such that $QN \perp PR$. If $PN \cdot NR = QN^2$, then prove that $\angle PQR = 90^\circ$.

Solution:

Given ΔPQR , N is a point on PR , such that $QN \perp PR$

and

$$PN \cdot NR = QN^2$$

To prove

$$\angle PQR = 90^\circ$$

Proof We have,

$$PN \cdot NR = QN^2$$

$\Rightarrow$

$$PN \cdot NR = QN \cdot QN$$

$\Rightarrow$

$$\frac{PN}{QN} = \frac{QN}{NR} \quad \dots(i)$$

In ΔQNP and ΔRNQ ,

$$\frac{PN}{QN} = \frac{QN}{NR}$$

and

$$\angle PNQ = \angle RNQ$$

$\therefore$

$$\Delta QNP \sim \Delta RNQ$$

Then, ΔQNP and ΔRNQ are equiangulars.

i.e.,

$$\angle PQN = \angle QRN$$

$$\angle RQN = \angle QPN$$

On adding both sides, we get

$$\angle PQN + \angle RQN = \angle QRN + \angle QPN$$

$\Rightarrow$

$$\angle PQR = \angle QRN + \angle QPN \quad \dots(ii)$$

We know that, sum of angles of a triangle = 180°

In ΔPQR , $\angle PQR + \angle QPR + \angle QRP = 180^\circ$

$\Rightarrow$

$$\angle PQR + \angle QPN + \angle QRN = 180^\circ \quad [\because \angle QPR = \angle QPN \text{ and } \angle QRP = \angle QRN]$$

$\Rightarrow$

$$\angle PQR + \angle PQR = 180^\circ \quad [\text{using Eq. (ii)}]$$

$\Rightarrow$

$$2 \angle PQR = 180^\circ$$


$\Rightarrow$

$$\angle PQR = \frac{180^\circ}{2} = 90^\circ$$

$\therefore$

$$\angle PQR = 90^\circ$$

Hence proved.



Question 12:

Areas of two similar triangles are 36 cm^2 and 100 cm^2 . If the length of a side of the larger triangle is 20 cm. Find the length of the corresponding side of the smaller triangle.

Solution:

Given, area of smaller triangle = 36 cm^2 and area of larger triangle = 100 cm^2

Also, length of a side of the larger triangle = 20 cm

Let length of the corresponding side of the smaller triangle = x cm

By property of area of similar triangle,

$$\begin{aligned}
 \Rightarrow & \frac{\text{ar (larger triangle)}}{\text{ar (smaller triangle)}} = \frac{(\text{Side of larger triangle})^2}{(\text{Side of smaller triangle})^2} \\
 \Rightarrow & \frac{100}{36} = \frac{(20)^2}{x^2} \Rightarrow x^2 = \frac{(20)^2 \times 36}{100} \\
 \Rightarrow & x^2 = \frac{400 \times 36}{100} = 144 \\
 \therefore & x = \sqrt{144} = 12 \text{ cm}
 \end{aligned}$$

Hence, the length of corresponding side of the smaller triangle is 12 cm.

Question 13:

In given figure, if $\angle ACB = \angle CDA$, $AC = 8$ cm and $AD = 3$ cm, then find BD .

Solution:


Given, $AC = 8$ cm, $AD = 3$ cm and

$$\angle ACB = \angle CDA$$

From figure,

$$\angle CDA = 90^\circ$$

$$\angle ACB = \angle CDA = 90^\circ$$



In right angled $\triangle ADC$,

$\Rightarrow$

$$AC^2 = AD^2 + CD^2$$

$\Rightarrow$

$$(8)^2 = (3)^2 + (CD)^2$$

$\Rightarrow$

$$64 - 9 = CD^2$$

In $\triangle CDB$ and $\triangle ADC$,

$$CD = \sqrt{55} \text{ cm}$$

$$\angle BDC = \angle ADC$$

[each 90°]

$$\angle DBC = \angle DCA$$

[each equal to $90^\circ - \angle A$]

$\therefore$

$$\triangle CDB \sim \triangle ADC$$

Then,

$$\frac{CD}{BD} = \frac{AD}{CD}$$

$\Rightarrow$

$$CD^2 = AD \times BD$$

$\therefore$

$$BD = \frac{CD^2}{AD} = \frac{(\sqrt{55})^2}{3} = \frac{55}{3} \text{ cm}$$

Question 14:



A 15 high tower casts a shadow 24 Long at a certain time and at the same time, a telephone pole casts a shadow 16 long. Find the height of the telephone pole.

Solution:

Let $BC = 15$ m be the tower and its shadow AB is 24 m. At that time $\angle CAB = \theta$. Again, let

$EF = h$ be a telephone pole and its shadow $DE = 16$ m. At the same time $\angle EDF = \theta$. Here,

$\triangle ABC$ and $\triangle DEF$ both are right angled triangles.



In $\triangle ABC$ and $\triangle DEF$,

$$\angle CAB = \angle EDF = \theta$$

$$\angle B = \angle E$$

[each 90°]

$\therefore$

$$\triangle ABC \sim \triangle DEF$$

[by AAA similarity criterion]

Then,

$$\frac{AB}{DE} = \frac{BC}{EF}$$

$\Rightarrow$

$$\frac{24}{16} = \frac{15}{h}$$

$\therefore$

$$h = \frac{15 \times 16}{24} = 10$$


Hence, the height of the telephone pole is 10 m.

Question 15:

Foot of a 10 m long ladder leaning against a vertical wall is 6 m away from the base of the wall. Find the height of the point on the wall where the top of the ladder reaches.

Solution:

Let AB be a vertical wall and AC = 10 m is a ladder. The top of the ladder reaches to A and distance of ladder from the base of the wall BC is 6 m.




$$\begin{aligned}
 &\text{In right angled } \triangle ABC, & AC^2 &= AB^2 + BC^2 & \text{[by Pythagoras theorem]} \\
 \Rightarrow & & (10)^2 &= AB^2 + (6)^2 \\
 \Rightarrow & & 100 &= AB^2 + 36 \\
 \Rightarrow & & AB^2 &= 100 - 36 = 64 \\
 \therefore & & AB &= \sqrt{64} = 8 \text{ cm}
 \end{aligned}$$

Hence, the height of the point on the wall where the top of the ladder reaches is 8 cm.

Exercise 6.4 Long Answer Type Questions

Question 1:

In given figure, if $\angle A = \angle C$, AB = 6 cm, BP = 15 cm, AP = 12 cm and CP = 4 cm, then find the lengths of PD and CD.



Solution:

Given, $\angle A = \angle C$, AB = 6 cm, BP = 15 cm, AP = 12 cm and CP = 4 cm

In $\triangle APB$ and $\triangle CPD$, $\angle A = \angle C$

[given]

$\angle APS = \angle CPD$

[vertically opposite angles]

$\therefore$

$\triangle APD \sim \triangle CPD$

[by AAA similarity criterion]

$\Rightarrow$

$$\frac{AP}{CP} = \frac{PB}{PD} = \frac{AB}{CD}$$

$\Rightarrow$

$$\frac{12}{4} = \frac{15}{PD} = \frac{6}{CD}$$

On taking first two terms, we get

$$\frac{12}{4} = \frac{15}{PD}$$

$$\Rightarrow PD = \frac{15 \times 4}{12} = 5 \text{ cm}$$

On taking first and last term, we get

$$\frac{12}{4} = \frac{6}{CD}$$

$$\Rightarrow CD = \frac{6 \times 4}{12} = 2 \text{ cm}$$

Hence, length of PD = 5 cm and length of CD = 2 cm

Question 2:

It is given that $\triangle ABC \sim \triangle EDF$ such that AB = 5 cm, AC = 7 cm, DF = 15 cm and DE = 12 cm. Find the lengths of the remaining sides of the triangles,


Solution:

Given, $\triangle ABC \sim \triangle EDF$, so the corresponding sides of $\triangle ABC$ and $\triangle EDF$ are in the same ratio.

i.e.,

$$\frac{AB}{ED} = \frac{AC}{EF} = \frac{BC}{DF}$$

...(i)



Also,

$$AB = 5 \text{ cm}, AC = 7 \text{ cm}$$

$$DF = 15 \text{ cm and } DE = 12 \text{ cm}$$

On putting these values in Eq. (i), we get

$$\frac{5}{12} = \frac{7}{EF} = \frac{BC}{15}$$

On taking first and second terms, we get

$$\frac{5}{12} = \frac{7}{EF}$$

⇒

$$EF = \frac{7 \times 12}{5} = 16.8 \text{ cm}$$

On taking first and third terms, we get

$$\frac{5}{12} = \frac{BC}{15}$$

⇒

$$BC = \frac{5 \times 15}{12} = 6.25 \text{ cm}$$

Hence, lengths of the remaining sides of the triangles are $EF = 16.8 \text{ cm}$ and $SC = 6.25 \text{ cm}$.

Question 3:


Prove that, if a line is drawn parallel to one side of a triangle to intersect the other two sides, then the two sides are divided in the same ratio.

Solution:

Let a $\triangle ABC$ in which a line DE parallel to BC intersects AB at D and AC at E . To prove DE divides the two sides in the same ratio.

i.e.,

$$\frac{AD}{DB} = \frac{AE}{EC}$$



Construction Join BE , CD and draw $EF \perp AB$ and $DG \perp AC$.

Proof Here,

$$\frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle BDE)} = \frac{\frac{1}{2} \times AD \times EF}{\frac{1}{2} \times DB \times EF} \quad [\because \text{area of triangle} = \frac{1}{2} \times \text{base} \times \text{height}]$$

$$= \frac{AD}{DB} \quad \dots (i)$$

similarly,

$$\frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle DEC)} = \frac{\frac{1}{2} \times AE \times DG}{\frac{1}{2} \times EC \times DG} = \frac{AE}{EC} \quad \dots (ii)$$

Now, since, $\triangle BDE$ and $\triangle DEC$ lie between the same parallel DE and BC and on the same base DE .

So,

$$\text{ar}(\triangle BDE) = \text{ar}(\triangle DEC) \quad \dots (iii)$$

From Eqs. (i), (ii) and (iii),

$$\frac{AD}{DB} = \frac{AE}{EC}$$


Hence proved.

Question 4:

In the given figure, if PQRS is a parallelogram and $AB \parallel PS$, then prove that $OC \parallel SR$.

Solution:

Given PQRS is a parallelogram, so $PQ \parallel SR$ and $PS \parallel QR$. Also, $AB \parallel PS$



To prove $OC \parallel SR$

Proof in $\triangle OPS$ and $\triangle OAB$,

$\therefore$

Then,

In $\triangle CQR$ and $\triangle CAB$,

$\therefore$

Then,

$\Rightarrow$

From Eqs. (i) and (ii),

On subtracting from both sides, we get

$\Rightarrow$

$\Rightarrow$

By converse of basic proportionality theorem,

$$PS \parallel AB$$

$$\angle POS = \angle AOB$$

$$\angle OSP = \angle OBA$$

$$\triangle OPS \sim \triangle OAB$$

$$\frac{PS}{AB} = \frac{OS}{OB}$$

$$QR \parallel PS \parallel AB$$

$$\angle QCR = \angle ACB$$

$$\angle CRQ = \angle CBA$$

$$\triangle CQR \sim \triangle CAB$$

$$\frac{QR}{AB} = \frac{CR}{CB}$$

$$\frac{PS}{AB} = \frac{CR}{CB}$$

$$\frac{OS}{OB} = \frac{CR}{CB} \text{ or } \frac{OB}{OS} = \frac{CB}{CR}$$

$$\frac{OB}{OS} - 1 = \frac{CB}{CR} - 1$$

$$\frac{OB - OS}{OS} = \frac{CB - CR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

[common angle]

[corresponding angles]

[by AAA similarity criterion]

... (i)

[common angle]

[corresponding angles]

... (ii)

[since, PQRS is a parallelogram, so $PS = QR$]

$$\frac{OS}{OB} = \frac{CR}{CB} \text{ or } \frac{OB}{OS} = \frac{CB}{CR}$$

$$\frac{OB}{OS} - 1 = \frac{CB}{CR} - 1$$

$$\frac{OB - OS}{OS} = \frac{CB - CR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$\frac{BS}{OS} = \frac{BR}{CR}$$

$$SR \parallel OC$$

Hence proved.

Question 5:

A 5 m long ladder is placed leaning towards a vertical wall such that it reaches the wall at a point 4 m high. If the foot of the ladder is moved 1.6 m towards the wall, then find the distance by which the top of the ladder would slide upwards on the wall.

Solution:

Let AC be the ladder of length 5 m and BC = 4 m be the height of the wall, which ladder is placed. If the foot of the ladder is moved 1.6 m towards the wall i.e., AD = 1.6 m, then the ladder is slide upward i.e., CE = x m.


In right angled $\triangle ABC$,

$$AC^2 = AB^2 + BC^2 \quad [\text{by Pythagoras theorem}]$$

$$\Rightarrow (5)^2 = (AB)^2 + (4)^2$$

$$\Rightarrow AB^2 = 25 - 16 = 9 \Rightarrow AB = 3 \text{ m}$$

$$\therefore DB = AB - AD = 3 - 1.6 = 1.4 \text{ m}$$



In right angled $\triangle EBD$,

$$\Rightarrow ED^2 = EB^2 + BD^2 \quad [\text{by Pythagoras theorem}]$$

$$\Rightarrow (5)^2 = (EB)^2 + (1.4)^2 \quad [\because BD = 1.4 \text{ m}]$$

$$\Rightarrow 25 = (EB)^2 + 1.96$$

$$\Rightarrow (EB)^2 = 25 - 1.96 = 23.04$$

$$\Rightarrow EB = \sqrt{23.04} = 4.8$$

Now, $EC = EB - BC = 4.8 - 4 = 0.8$

Hence, the top of the ladder would slide upwards on the wall at distance 0.8 m.

Question 6:

For going to a city B from city A there is a route via city C such that $AC \perp CB$, $AC = 2x$ km and $CB = 2(x + 7)$ km. It is proposed to construct a 26 km highway which directly connects the two cities A and B. Find how much distance will be saved in reaching city B from city A after the construction of the highway.

Solution:

Given, $AC \perp CB$, km, $CB = 2(x + 7)$ km and $AB = 26$ km

On drawing the figure, we get the right angled $\triangle ACB$ right angled at C.

Now, In $\triangle ACB$, by Pythagoras theorem,

$$\Rightarrow AB^2 = AC^2 + BC^2$$

$$\Rightarrow (26)^2 = (2x)^2 + \{2(x + 7)\}^2$$

$$\Rightarrow 676 = 4x^2 + 4(x^2 + 49 + 14x)$$

$$\Rightarrow 676 = 4x^2 + 4x^2 + 196 + 56x$$

$$\Rightarrow 676 = 8x^2 + 56x + 196$$

$$\Rightarrow 8x^2 + 56x - 480 = 0$$

$$\Rightarrow \text{On dividing by 8, we get } x^2 + 7x - 60 = 0$$

$$\Rightarrow x^2 + 12x - 5x - 60 = 0$$

$$\Rightarrow x(x + 12) - 5(x + 12) = 0$$

$$\Rightarrow (x + 12)(x - 5) = 0$$

$$\therefore x = -12, x = 5$$

Since, distance cannot be negative.

$$\therefore x = 5 \quad [\because x \neq -12]$$

Now,

$$AC = 2x = 10 \text{ km}$$

and

$$BC = 2(x + 7) = 2(5 + 7) = 24 \text{ km}$$

The distance covered to reach city B from city A via city C

$$= AC + BC$$


$$= 10 + 24$$

$$= 34 \text{ km}$$

Distance covered to reach city B from city A after the construction of the highway

$$= BA = 26 \text{ km}$$

Hence, the required saved distance is $34 - 26$ i.e., 8 km.




Question 7:

A flag pole 18 m high casts a shadow 9.6 m long. Find the distance of the top of the pole from the far end of the shadow.

Solution:

Let $BC = 18$ m be the flag pole and its shadow be $AB = 9.6$ m. The distance of the top of the pole, C from the far end i.e., A of the shadow is AC .



In right angled $\triangle ABC$,

$\Rightarrow$

$\Rightarrow$

$\therefore$

$$AC^2 = AB^2 + BC^2$$

[by Pythagoras theorem]

$$AC^2 = (9.6)^2 + (18)^2$$

$$AC^2 = 92.16 + 324$$

$$AC^2 = 416.16$$

$$AC = \sqrt{416.16} = 20.4\text{m}$$


Hence, the required distance is 20.4 m.

Question 8:

A street light bulb is fixed on a pole 6 m above the level of the street. If a woman of height 1.5 m casts a shadow of 3 m, then find how far she is away from the base of the pole.

Solution:

Let A be the position of the street bulb fixed on a pole $AB = 6$ m and $CD = 1.5$ m be the height of a woman and her shadow be $ED = 3$ m. Let distance between pole and woman be x m.



Here, woman and pole both are standing vertically.

So,

In $\triangle CDE$ and $\triangle ABE$,

$\therefore$

Then,

$\Rightarrow$

$\Rightarrow$

$\Rightarrow$

$\Rightarrow$

$\therefore$

$$CD \parallel AB$$

$$\angle E = \angle E$$

$$\angle ABE = \angle CDE$$

$$\triangle CDE \sim \triangle ABE$$

$$\frac{ED}{EB} = \frac{CD}{AB}$$

$$\frac{3}{3+x} = \frac{1.5}{6}$$

$$3 \times 6 = 1.5(3+x)$$

$$18 = 1.5 \times 3 + 1.5x$$

$$1.5x = 18 - 4.5$$

$$x = \frac{13.5}{1.5} = 9\text{m}$$

[common angle]


[each equal to 90°]

[by AAA similarity criterion]

Hence, she is at the distance of 9 m from the base of the pole.

Question 9:

In given figure, ABC is a triangle right angled at B and $BD \perp AC$. If $AD = 4$ cm and $CD = 5$ cm, then find BD and AB .



Solution:

Given, $\triangle ABC$ in which $\angle B = 90^\circ$ and $BD \perp AC$

Also, $AD = 4 \text{ cm}$ and $CD = 5 \text{ cm}$

In $\triangle ADB$ and $\triangle CDB$,

$$\angle ADB = \angle CDB$$

[each equal to 90°]

and

$$\angle BAD = \angle DCB$$

[each equal to $90^\circ - \angle C$]

$\therefore$

$$\triangle DBA \sim \triangle DCB$$

[by AAA similarity criterion]

Then,

$$\frac{DB}{DA} = \frac{DC}{DB}$$

$\Rightarrow$

$$DB^2 = DA \times DC$$

$\Rightarrow$

$$DB^2 = 4 \times 5$$

$\Rightarrow$

$$DB = 2\sqrt{5} \text{ cm}$$

In right angled $\triangle BDC$,

$$BC^2 = BD^2 + CD^2$$

[by Pythagoras theorem]

$\Rightarrow$

$$BC^2 = (2\sqrt{5})^2 + (5)^2$$

$$= 20 + 25 = 45$$

$\Rightarrow$

$$BC = \sqrt{45} = 3\sqrt{5}$$

Again,

$$\triangle DBA \sim \triangle DCB,$$

$\therefore$

$$\frac{DB}{DC} = \frac{BA}{BC}$$

$\Rightarrow$

$$\frac{2\sqrt{5}}{5} = \frac{BA}{3\sqrt{5}}$$


$\therefore$

$$BA = \frac{2\sqrt{5} \times 3\sqrt{5}}{5} = 6 \text{ cm}$$

Hence, $BD = 2\sqrt{5} \text{ cm}$ and $AB = 6 \text{ cm}$

Question 10:

In given figure PQR is a right triangle, right angled at Q and $QS \perp PR$. If $PQ = 6 \text{ cm}$ and $PS = 4 \text{ cm}$, then find QS, RS and QR.



Solution:

Given, $\triangle PQR$ in which $\angle Q = 90^\circ$, $QS \perp PR$ and $PQ = 6 \text{ cm}$, $PS = 4 \text{ cm}$ In $\triangle SQP$ and $\triangle SRQ$,

$$\angle PSQ = \angle RSQ$$

[each equal to 90°]

$$\angle SPQ = \angle SQR$$

[each equal to $90^\circ - \angle R$]

$\therefore$

$$\triangle SQP \sim \triangle SRQ$$

Then,

$$\frac{SQ}{PS} = \frac{SR}{SQ}$$

$\Rightarrow$

$$SQ^2 = PS \times SR$$

...(i)

In right angled ΔPSQ , $PQ^2 = PS^2 + QS^2$ [by Pythagoras theorem]
 $\Rightarrow (6)^2 = (4)^2 + QS^2$
 $\Rightarrow 36 = 16 + QS^2$
 $\Rightarrow QS^2 = 36 - 16 = 20$
 $\therefore QS = \sqrt{20} = 2\sqrt{5} \text{ cm}$
 On putting the value of QS in Eq. (i), we get
 $(2\sqrt{5})^2 = 4 \times SR$
 $\Rightarrow SR = \frac{4 \times 5}{4} = 5 \text{ cm}$
 In right angled ΔQSR , $QR^2 = QS^2 + SR^2$
 $\Rightarrow QR^2 = (2\sqrt{5})^2 + (5)^2$
 $\Rightarrow QR^2 = 20 + 25$
 $\therefore QR = \sqrt{45} = 3\sqrt{5} \text{ cm}$
 Hence, $QS = 2\sqrt{5} \text{ cm}$, $RS = 5 \text{ cm}$ and $QR = 3\sqrt{5} \text{ cm}$

Question 11:

In ΔPQR , $PD \perp QR$ such that D lies on QR, if $PQ = a$, $PR = b$, $QD = c$ and $DR = d$, then prove that $(a + b)(a - b) = (c + d)(c - d)$.


Solution:

Given In ΔPQR , $PD \perp QR$, $PQ = a$, $PR = b$, $QD = c$ and $DR = d$

To prove $(a + b)(a - b) = (c + d)(c - d)$

Proof In right angled ΔPDQ ,

$PQ^2 = PD^2 + QD^2$ [by Pythagoras theorem]
 $\Rightarrow a^2 = PD^2 + c^2$
 $\Rightarrow PD^2 = a^2 - c^2$... (i)



In right angled ΔPDR , $PR^2 = PD^2 + DR^2$ [by Pythagoras theorem]
 $\Rightarrow b^2 = PD^2 + d^2$
 $\Rightarrow PD^2 = b^2 - d^2$... (ii)

From Eqs. (i) and (ii),

$a^2 - c^2 = b^2 - d^2$
 $\Rightarrow a^2 - b^2 = c^2 - d^2$
 $\Rightarrow (a - b)(a + b) = (c - d)(c + d)$ Hence proved.

Question 12:

In a quadrilateral ΔBCD , $\angle A + \angle D = 90^\circ$. Prove that

$$AC^2 + BD^2 = AD^2 + BC^2.$$

Solution:

Given Quadrilateral ΔBCD , in which $\angle A + \angle D = 90^\circ$

To prove

$$AC^2 + BD^2 = AD^2 + BC^2$$

Construct Produce AB and CD to meet at E.

Also, join AC and BD.

Proof In $\triangle AED$, $\angle A + \angle D = 90^\circ$ [given]
 $\therefore \angle E = 180^\circ - (\angle A + \angle D) = 90^\circ$
 [∵ sum of angles of a triangle = 180°]

Then, by Pythagoras theorem, $AD^2 = AE^2 + DE^2$

In $\triangle BEC$, by Pythagoras theorem, $BC^2 = BE^2 + EC^2$

On adding both equations, we get

$$AD^2 + BC^2 = AE^2 + DE^2 + BE^2 + EC^2 \quad \dots(i)$$

In $\triangle AEC$, by Pythagoras theorem,

$$AC^2 = AE^2 + EC^2$$

and in $\triangle BED$, by Pythagoras theorem,

$$BD^2 = BE^2 + DE^2$$


On adding both equations, we get

$$AC^2 + BD^2 = AE^2 + EC^2 + BE^2 + DE^2 \quad \dots(ii)$$

From Eqs. (i) and (ii),


$$AC^2 + BD^2 = AD^2 + BC^2$$

Hence proved.



Question 13:

In given figure, $l \parallel m$ and line segments AB, CD and EF are concurrent at point P. Prove that $\frac{AE}{BF} = \frac{AC}{BD} = \frac{CE}{FD}$.



Solution:

Given $l \parallel m$ and line segments AB, CD and EF are concurrent at point P.

To prove

$$\frac{AE}{BF} = \frac{AC}{BD} = \frac{CE}{FD}$$

Proof In $\triangle APC$ and $\triangle BPD$,

$$\angle APC = \angle BPD$$

[vertically opposite angles]

$$\angle PAC = \angle PBD$$

[alternate angles]

$\therefore$

$$\triangle APC \sim \triangle BPD$$

[by AAA similarity criterion]

Then,

$$\frac{AP}{PB} = \frac{AC}{BD} = \frac{PC}{PD} \quad \dots(i)$$

In $\triangle APE$ and $\triangle BPF$,

$$\angle APE = \angle BPF$$

[vertically opposite angles]

$$\angle PAE = \angle PBF$$

[alternate angles]

$\therefore$

$$\triangle APE \sim \triangle BPF$$

[by AAA similarity criterion]

Then,

$$\frac{AP}{PB} = \frac{AE}{BF} = \frac{PE}{PF} \quad \dots(ii)$$

In $\triangle PEC$ and $\triangle PFD$,

$$\angle EPC = \angle FPD$$

[vertically opposite angles]

$$\angle PCE = \angle PFD$$

[alternate angles]

$\therefore$

$$\triangle PEC \sim \triangle PFD$$

[by AAA similarity criterion]

Then,

$$\frac{PE}{PF} = \frac{PC}{PD} = \frac{EC}{FD} \quad \dots(iii)$$

From Eqs. (i), (ii) and (iii),

$$\frac{AP}{PB} = \frac{AC}{BD} = \frac{AE}{BF} = \frac{PE}{PF} = \frac{EC}{FD}$$

$\therefore$


$$\frac{AE}{BF} = \frac{AC}{BD} = \frac{CE}{FD}$$

Hence proved.

Question 14:

In figure, PA, QB, RC and SD are all perpendiculars to a line l , AB = 6 cm, BC = 9 cm, CD =

12 cm and $SP = 36$ cm. Find PQ , QR and RS .



Solution:

Given, $AS = 6$ cm, $BC = 9$ cm, $CD = 12$ cm and $SP = 36$ cm

Also, PA , QB , RC and SD are all perpendiculars to line l .

$PA \parallel QS \parallel SC \parallel SD$

By basic proportionality theorem,

$$PQ : QR : RS = AB : BC : CD \\ = 6 : 9 : 12$$

$$PQ = 6x, QR = 9x \text{ and } RS = 12x$$

$$PS = 36 \text{ km}$$

$$PQ + QR + RS = 36$$

$$6x + 9x + 12x = 36$$

$$27x = 36$$

$$x = \frac{36}{27} = \frac{4}{3}$$

$$PQ = 6x = 6 \times \frac{4}{3} = 8 \text{ cm}$$

$$QR = 9x = 9 \times \frac{4}{3} = 12 \text{ cm}$$

$$RS = 12x = 12 \times \frac{4}{3} = 16 \text{ cm}$$

Question 15:

O is the point of intersection of the diagonals AC and BD of a trapezium $ABCD$ with $AB \parallel DC$.

Through O , a line segment PQ is drawn parallel to AB meeting AD in P and BC in Q , prove that $PO = QO$.


Solution:

Given $ABCD$ is a trapezium. Diagonals AC and BD are intersect at O .

$PQ \parallel AB \parallel DC$.

To prove

$$PO = QO$$



Proof In $\triangle ABD$ and $\triangle POD$,

$$PO \parallel AB$$

$$\angle D = \angle D$$

$$\angle ABD = \angle POD$$

$$\triangle ABD \sim \triangle POD$$

$$\frac{OP}{AB} = \frac{PD}{AD}$$

$\therefore$

Then,

In $\triangle ABC$ and $\triangle OQC$,

$$OQ \parallel AB$$

$$\angle C = \angle C$$

$$\angle BAC = \angle QOC$$

$$\triangle ABC \sim \triangle OQC$$

$\therefore$

Then,

$$\frac{OQ}{AB} = \frac{QC}{BC}$$

$$[\because PQ \parallel AB]$$

[common angle]

[corresponding angles]

[by AAA similarity criterion]

...(i)

$$[\because OQ \parallel AB]$$

[common angle]

[corresponding angle]

[by AAA similarity criterion]

...(ii)

Now, in $\triangle ADC$,

$\therefore$

In $\triangle ABC$,

$\therefore$

From Eqs. (iii) and (iv),

$\therefore$

Adding 1 on both sides, we get

$\Rightarrow$

$\Rightarrow$

$\Rightarrow$

$\Rightarrow$

$\Rightarrow$

$\Rightarrow$

$$OP \parallel DC$$

$$\frac{AP}{PP} = \frac{OA}{OC}$$

[by basic proportionality theorem]... (iii)

$$OQ \parallel AB$$

$$\frac{BQ}{QC} = \frac{OA}{OC}$$

[by basic proportionality theorem]... (iv)

$$\frac{AP}{PD} = \frac{BQ}{QC}$$

$$\frac{AP}{PD} + 1 = \frac{BQ}{QC} + 1$$

$$\frac{AP + PD}{PD} = \frac{BQ + QC}{QC}$$

$$\frac{AD}{PD} = \frac{BC}{QC}$$

$$\frac{AD}{PD} = \frac{BC}{QC}$$

$$\frac{AD}{PD} = \frac{BC}{QC}$$

$$\frac{AD}{OP} = \frac{BC}{OQ}$$

$$\frac{AB}{OP} = \frac{BC}{OQ}$$

$$\frac{AB}{OP} = \frac{BC}{OQ}$$

$$\frac{AB}{AB} = \frac{OQ}{OQ}$$

$$OP = OQ$$

[from Eqs. (i) and (ii)]


[from Eq. (ii)]

Hence proved.

Question 16:

In figure, line segment DF intersects the side AC of a $\triangle ABC$ at the point E such that E is the mid-point of CA and $\angle AEF = \angle AFE$. Prove that $\frac{BD}{CD} = \frac{BF}{CF}$.

$\angle AEF = \angle AFE$. Prove that $\frac{BD}{CD} = \frac{BF}{CF}$.



Solution:

Given $\triangle ABC$, E is the mid-point of CA and $\angle AEF = \angle AFE$

To prove

$$\frac{BD}{CD} = \frac{BF}{CF}$$

Construction Take a point G on AB such that $CG \parallel EF$.

Proof Since, E is the mid-point of CA,

$\therefore$

$$CE = AE$$

... (i)

In $\triangle ACG$, $CG \parallel EF$ and E is mid-point of CA.

So,

$$CE = GF$$

... (ii)

[by mid-point theorem]

Now, in $\triangle BCG$ and $\triangle BDF$,

$\therefore$

$$CG \parallel EF$$

$$\frac{BC}{CD} = \frac{BG}{GF}$$

[by basic proportionality theorem]

$\Rightarrow$

$$\frac{BC}{CD} = \frac{BG}{GF}$$

$$\frac{BC}{CD} = \frac{BF - GF}{GF} \Rightarrow \frac{BC}{CD} = \frac{BF}{GF} - 1$$

$\Rightarrow$

$$\frac{BC}{CD} + 1 = \frac{BF}{GF}$$

[from Eq. (ii)]


$\Rightarrow$

$$\frac{BC + CD}{CD} = \frac{BF}{GF}$$

$$\frac{BD}{CD} = \frac{BF}{GF}$$

$$\frac{BD}{CD} = \frac{BF}{CF}$$

Hence proved.



Question 17:

Prove that the area of the semi-circle drawn on the hypotenuse of a right angled triangle is equal to the sum of the areas of the semi-circles drawn on the other two sides of the triangle.

Solution:

Let ABC be a right triangle, right angled at B and AB = y, BC = x.

Three semi-circles are drawn on the sides AB, BC and AC, respectively with diameters AB, BC and AC, respectively.

Again, let area of circles with diameters AB, BC and AC are respectively A_1 , A_2 and A_3 .

To prove $A_3 = A_1 + A_2$

Proof In $\triangle ABC$, by Pythagoras theorem,

$$AC^2 = AB^2 + BC^2$$

$$\Rightarrow AC^2 = y^2 + x^2$$

$$\Rightarrow AC = \sqrt{y^2 + x^2}$$

We know that, area of a semi-circle with radius, $r = \frac{\pi r^2}{2}$

$$\therefore \text{Area of semi-circle drawn on AC, } A_3 = \frac{\pi}{2} \left(\frac{AC}{2} \right)^2 = \frac{\pi}{2} \left(\frac{\sqrt{y^2 + x^2}}{2} \right)^2$$



$$A_3 = \frac{\pi(y^2 + x^2)}{8} \quad \dots(i)$$

Now, area of semi-circle drawn on AB, $A_1 = \frac{\pi}{2} \left(\frac{AB}{2} \right)^2$

$$\Rightarrow A_1 = \frac{\pi}{2} \left(\frac{y}{2} \right)^2 \Rightarrow A_1 = \frac{\pi y^2}{8} \quad \dots(ii)$$


and area of semi-circle drawn on BC, $A_2 = \frac{\pi}{2} \left(\frac{BC}{2} \right)^2 = \frac{\pi}{2} \left(\frac{x}{2} \right)^2$

$$\Rightarrow A_2 = \frac{\pi x^2}{8}$$

On adding Eqs. (ii) and (iii), we get $A_1 + A_2 = \frac{\pi y^2}{8} + \frac{\pi x^2}{8}$

$$= \frac{\pi(y^2 + x^2)}{8} = A_3 \quad [\text{from Eq. (i)}]$$

$\Rightarrow A_1 + A_2 = A_3$ **Hence proved.**



Question 18:

Prove that the area of the equilateral triangle drawn on the hypotenuse of a right angled triangle is equal to the sum of the areas of the equilateral triangle drawn on the other two sides of the triangle.

Solution:

Let a right triangle BAC in which $\angle A$ is right angle and AC = y, AB = x.

Three equilateral triangles $\triangle AEC$, $\triangle AFB$ and $\triangle CBD$ are drawn on the three sides of $\triangle ABC$.

Again let area of triangles made on AC, AS and BC are A_1 , A_2 and A_3 , respectively.

To prove $A_3 = A_1 + A_2$

Proof In $\triangle CAB$, by Pythagoras theorem,

$$BC^2 = AC^2 + AB^2$$

$$\Rightarrow BC^2 = y^2 + x^2$$

$$\Rightarrow BC = \sqrt{y^2 + x^2}$$

We know that, area of an equilateral triangle = $\frac{\sqrt{3}}{4} (\text{Side})^2$


$$\therefore \text{Area of equilateral } \triangle AEC, A_1 = \frac{\sqrt{3}}{4} (AC)^2$$

$$\Rightarrow A_1 = \frac{\sqrt{3}}{4} y^2 \quad \dots(i)$$

and area of equilateral $\triangle AFB$, $A_2 = \frac{\sqrt{3}}{4} (AB)^2 = \frac{\sqrt{3}}{4} \sqrt{y^2 + x^2}$

$$= \frac{\sqrt{3}}{4} (y^2 + x^2) = \frac{\sqrt{3}}{4} y^2 + \frac{\sqrt{3}}{4} x^2$$

$$= A_1 + A_2$$



[from Eqs. (i) and (ii)]
Hence proved.